

Piotr Achinger

Introduction to non-Archimedean Geometry

NON-ARCHIMEDEAN or rigid-analytic geometry is an analog of complex analytic geometry over non-Archimedean fields, such as the field of p -adic numbers \mathbb{Q}_p or the field of formal Laurent series $k((t))$. It was introduced and formalized by Tate in the 1960s, whose goal was to understand elliptic curves over a p -adic field by means of a uniformization similar to the familiar description of an elliptic curve over \mathbb{C} as quotient of the complex plane by a lattice. It has since gained status of a foundational tool in algebraic and arithmetic geometry, and several other approaches have been found by Raynaud, Berkovich, and Huber. In recent years, it has become even more prominent with the work of Scholze and Kedlaya in p -adic Hodge theory, as well as the non-Archimedean approach to mirror symmetry proposed by Kontsevich. That said, we still do not know much about rigid-analytic varieties, and many foundational questions remain unanswered.

The goal of this lecture course is to introduce the basic notions of rigid-analytic geometry. We will assume familiarity with schemes.

Problem sets and other materials related to the course are available at

<http://achinger.impan.pl/lecture20f.html>

Our basic reference is the book *Lectures on Formal and Rigid Geometry* by Siegfried Bosch. More references are found in the text.

This document uses the *tufte-book* L^AT_EXdocument class based on the design of Edward Tufte's books; some typesetting tricks were shamelessly stolen from Eric Peterson's lecture notes.

These notes owe a lot to the generous help of Alex Youcis.

Contents

1	Two interpretations of non-Archimedean geometry	5
1.1	First example: the unit disc	7
1.2	Tate's admissible topology on the unit disc	7
1.3	Raynaud's approach	9
1.4	Why study rigid geometry?	10
2	Non-archimedean fields	13
2.1	Valuation rings and valuations	13
2.2	Valuations and norms	15
2.3	Geometric examples of valuations	16
2.4	Nonarchimedean fields	18
2.5	Extensions of nonarchimedean fields	20
2.6	Slopes of the Newton polygon	22
2.A	Henselian rings	23
3	The Tate algebra	27
3.1	Definition of the Tate algebra	27
3.2	The topology on $K\langle X_1, \dots, X_r \rangle$ and the Gauss norm	28
3.3	The universal property	30
3.4	The Tate algebra is Noetherian	31
3.5	Maximal ideals	33
3.6	More commutative algebra	34
3.A	Banach spaces (with Alex Youcis)	36
4	Affinoid algebras and spaces	43
4.1	Affinoid algebras and the residue norm	43
4.2	The supremum norm	43
4.3	The canonical topology	46
5	Sheaves, sites, and topoi	47
5.1	Motivation: reinventing the real	47
5.2	Sites	49
5.3	G -topologies	51
5.A	Sober topological spaces	53
6	Bibliography	57

1

Two interpretations of non-Archimedean geometry

THE p -ADIC NUMBERS \mathbf{Q}_p are usually defined either as the completion of the rational numbers \mathbf{Q} with respect to the p -adic absolute value

$$\left| \frac{a}{b} \right|_p = p^{\text{ord}_p b - \text{ord}_p a}, \quad (1.1)$$

or as the fraction field of the p -adic integers \mathbf{Z}_p defined as the inverse limit

$$\mathbf{Z}_p = \varprojlim_n \mathbf{Z}/p^n\mathbf{Z}. \quad (1.2)$$

We can refer to (1.1) as the “metric” or “analytic” point of view, while (1.2) represents a more “algebraic” (or “formal”) perspective.¹

Both interpretations have their advantages and drawbacks. The metric approach is admittedly closer to one’s intuition, and allows one to employ right away the powerful tools of topology and analysis. However, the topology of the p -adic numbers is quite pathological: \mathbf{Q}_p is a totally disconnected topological space. This makes it difficult to proceed by analogy with real or complex analysis.

The algebraic approach allows us to reduce questions about \mathbf{Q}_p to pure algebra over the rather simple rings $\mathbf{Z}/p^n\mathbf{Z}$. One therefore has commutative algebra and algebraic geometry at their disposal, which, in turn, allows one to more easily make sound and precise arguments. The downside: the relationship between objects over \mathbf{Q}_p and over $\mathbf{Z}/p^n\mathbf{Z}$ can often be extremely convoluted.

TO ACHIEVE p -ADIC ENLIGHTENMENT, one needs a good grasp of both², as well as a means of switching between the two with ease. The goal of these lectures is to explain how to do p -adic geometry (or, more generally, non-Archimedean geometry³) by combining the analytic and the algebraic approaches. Roughly speaking, the first will be represented by Tate’s notion of rigid analytic varieties, and the second by Raynaud’s approach using formal schemes.

WE WILL NOW GO BEYOND p -adic numbers and fix the notation which we will use most of the time. By a *non-Archimedean field* we mean a field K equipped with a non-Archimedean norm, which by definition is a function

$$|\cdot|: K \rightarrow [0, \infty)$$

¹ We choose to ignore here the (rather useless) definition of p -adic numbers in terms of base- p digit expansions.

² *It seems as though we must use sometimes the one theory and sometimes the other, while at times we may use either. We are faced with a new kind of difficulty. We have two contradictory pictures of reality; separately neither of them fully explains the phenomena of light, but together they do.*

A. Einstein, L. Infeld *The Evolution of Physics*

³ More precisely, *rigid (or rigid-analytic) geometry*, whose strange name we will justify later on.

such that

1. $|x| = 0$ if and only if $x = 0$,
2. $|xy| = |x| \cdot |y|$,
3. $|x + y| \leq \max(|x|, |y|)$.

We also assume that $|x| \neq 1$ for some $x \neq 0$ (i.e. that $|\cdot|$ is *nontrivial*), and that K is *complete* with respect to (the metric defined by) the norm.⁴

The third axiom, stronger than the triangle inequality $|x + y| \leq |x| + |y|$, is what makes the field non-Archimedean. It implies that the subset

$$\mathcal{O} = \{x \in K \text{ such that } |x| \leq 1\}$$

is a subring of K , called the *valuation ring*. It is local with maximal ideal

$$\mathfrak{m} = \{x \in K \text{ such that } |x| < 1\}.$$

We denote the residue field \mathcal{O}/\mathfrak{m} by k .

Let $t \in \mathfrak{m}$ be a nonzero element.⁵ Completeness of K is equivalent to the fact that the natural map

$$\mathcal{O} \rightarrow \varprojlim_n \mathcal{O}/t^n \mathcal{O}$$

is an isomorphism. The field K can be recovered as the fraction field of \mathcal{O} , in fact it is the localization $K = \mathcal{O}[\frac{1}{t}]$. The inverse limit above carries the inverse limit topology (with the $\mathcal{O}/t^n \mathcal{O}$ being equipped with the discrete topology), and the isomorphism is an isomorphism of topological rings if \mathcal{O} has the metric topology induced by the norm $|\cdot|$. The topology on K is the unique one with respect to which \mathcal{O} is an *open* subring. This implies that K is encoded as a *topological field* by the inverse system above.

The basic examples are complete discrete valuation fields (cdvf), which can be characterized as those K as above for which the maximal ideal \mathfrak{m} is principal, so that \mathcal{O} is a complete discrete valuation ring (cdvr) with maximal ideal \mathfrak{m} , residue field $k = \mathcal{O}/\mathfrak{m}$, and fraction field K . Naturally, our main example is

$$\mathcal{O} = \mathbf{Z}_p, \quad K = \mathbf{Q}_p, \quad \mathfrak{m} = (p), \quad k = \mathbf{F}_p,$$

and another one is the *Laurent series field* (over a base field k)⁶

$$\mathcal{O} = k[[t]] := \varprojlim_n k[t]/(t^n), \quad K = k((t)) := \mathcal{O}\left[\frac{1}{t}\right].$$

The characteristic of k is called the *residue characteristic* of K . If it is equal to the characteristic of K , we say that K is of *equal characteristic*, otherwise it is of *mixed characteristic*. In the latter case, K has characteristic zero. Thus \mathbf{Q}_p and its normed extensions are of mixed characteristic, and the fields $k((t))$ have equal characteristic. In fact, every cdvf of equal characteristic is of the form $k((t))$.

In general, we will have to work with non-Archimedean fields K which are not cdvf's, in which case the valuation ring \mathcal{O} is non-Noetherian. Indeed, it is often useful to consider K algebraically closed, while a complete discrete valuation field is never algebraically closed.⁷

⁴ In some sources, non-Archimedean fields are not assumed to be complete and/or nontrivially valued.

⁵ We call such a t a *pseudouniformizer*.

⁶ Intuition: $k((t))$ is the field of functions on the “infinitesimal punctured disc”

$$\text{Spec } k((t)) = \text{Spec } k[[t]] \setminus \{t = 0\}.$$

⁷ Consider a generator of \mathfrak{m} , i.e. an element of valuation one. Does it have a square root in K ?

1.1 First example: the unit disc

The study of schemes begins with the case of the affine line over a base field k

$$\mathbf{A}_k^1 = \text{Spec } k[x],$$

from which one obtains \mathbf{A}_k^n by direct product, then affine schemes of finite type over k by taking closed subschemes $X \subseteq \mathbf{A}_k^n$, and finally schemes locally of finite type over k by gluing. If k is algebraically closed, then by Hilbert's Nullstellensatz, closed points of \mathbf{A}_k^1 are in bijection with k .

In non-Archimedean geometry over an algebraically closed⁸ non-Archimedean field K , similar role is played by the closed unit disc

$$\mathbf{D}_K^1 = \{x \in K : |x| \leq 1\}.$$

Proceeding by analogy with scheme theory, we start with the algebra of functions on \mathbf{D}_K^1 , which should consist of power series $f = \sum_{n \geq 0} a_n x^n$ which converge for $|x| \leq 1$. An easy check shows that a series in K converges if and only if its terms tend to zero. We conclude that we want the ring of "holomorphic functions" on \mathbf{D}_K^1 to be

$$K\langle X \rangle = \left\{ \sum_{n \geq 0} a_n X^n \in K[[X]] \text{ with } a_n \rightarrow 0 \text{ as } n \rightarrow \infty \right\}.$$

Next, we would like to equip \mathbf{D}_K^1 with a *sheaf* of functions whose global sections is the above algebra $K\langle X \rangle$. The naive idea is to define, for an open subset $U \subseteq \mathbf{D}_K^1$, the ring $\mathcal{O}^{\text{wobbly}}(U)$ as the set of functions $U \rightarrow K$ which can be represented locally as a power series.

Indeed, this is trivially a sheaf, and we do obtain an injection

$$K\langle X \rangle \rightarrow \mathcal{O}^{\text{wobbly}}(\mathbf{D}_K^1).$$

However, this map is very far from being surjective. Indeed, \mathbf{D}_K^1 is highly disconnected, for example

$$\mathbf{D}_K^1 = \{|x| = 1\} \cup \{|x| < 1\} \tag{1.3}$$

expresses \mathbf{D}_K^1 as a union of two disjoint open (!) subsets. The function $f \in \mathcal{O}(\mathbf{D}_K^1)$ equal to 1 on the first open and 0 on the second is not in the image of $K\langle X \rangle$. (This example justifies the adjective *wobbly*.) Clearly, something goes terribly wrong with analytic continuation in the nonarchimedean setting!

1.2 Tate's admissible topology on the unit disc

The first attempt at fixing this issue is due to Krasner, and is based on a non-Archimedean analog of Runge's theorem in complex analysis. A *Krasner analytic function* on \mathbf{D}_K^1 is a uniform limit of rational functions with no poles inside \mathbf{D}_K^1 . This leads to a presheaf \mathcal{O} for which $\mathcal{O}(\mathbf{D}_K^1) = K\langle X \rangle$, and which has the property that $\mathcal{O}(U)$ is a domain if U "should be" connected. Still, it is not a sheaf.

⁸ We make this assumption only for simplicity and only in this introduction.

Let us explain, in a simple case, Tate’s idea of fixing the issue. Consider the following covering of \mathbf{D}_K^1 :

$$\mathbf{D}_K^1 = \underbrace{\{|x| \leq \rho\}}_U \cup \underbrace{\{\rho \leq |x| \leq 1\}}_V \tag{1.4}$$

with $0 < \rho < 1$, $\rho = |t|$ for some $t \in K$. The algebra of (Krasner analytic) functions $\mathcal{O}(U)$ on the smaller disc $U = \{|x| \leq \rho\}$ consists of power series converging on this disc, i.e.

$$K\left\langle \frac{X}{t} \right\rangle = \left\{ f = \sum_{n \geq 0} a_n X^n \in K[[X]] : \lim_{n \rightarrow \infty} |a_n| \rho^n = 0 \right\}.$$

Similarly, for the annulus $V = \{\rho \leq |x| \leq 1\}$, $\mathcal{O}(V)$ consists of convergent Laurent series

$$K\left\langle X, \frac{t}{X} \right\rangle = \left\{ f = \sum_{n \in \mathbf{Z}} a_n X^n : \lim_{n \rightarrow \infty} |a_n| = 0, \lim_{n \rightarrow -\infty} |a_n| \rho^n = 0 \right\},$$

and functions $\mathcal{O}(U \cap V)$ on the intersection $U \cap V = \{|x| = \rho\}$ are

$$K\left\langle \frac{X}{t}, \frac{t}{X} \right\rangle = \left\{ f = \sum_{n \in \mathbf{Z}} a_n X^n : \lim_{|n| \rightarrow \infty} |a_n| \rho^n = 0 \right\}.$$

It turns out that we are lucky: the sequence

$$0 \rightarrow K\langle X \rangle \rightarrow K\left\langle \frac{X}{t} \right\rangle \times K\left\langle X, \frac{t}{X} \right\rangle \rightarrow K\left\langle \frac{X}{t}, \frac{t}{X} \right\rangle \tag{1.5}$$

is exact.⁹ Thus \mathcal{O} satisfies the sheaf condition with respect to the covering $U \cup V$.

⁹ Check this!

TATE’S SOLUTION is now to identify a class of *admissible coverings* $U = \bigcup U_i$ of opens $U \subseteq \mathbf{D}_K^1$. For $U = \mathbf{D}_K^1$, these are the coverings admitting a *finite* refinement by subsets of the form

$$\{|x - a| \leq |t|, |x - a_i| \geq |t_i|\}.$$


The covering (1.3) is not admissible in this sense, while (1.4) is. *Tate’s acyclicity theorem* says that the presheaf \mathcal{O} satisfies the sheaf condition for all admissible coverings. Exactness of (1.5) is a basic special case.

In particular, this implies that \mathbf{D}_K^1 is *quasi-compact* with respect to the admissible topology: every *admissible* cover admits a finite subcover. Moreover, it becomes *connected* in the sense that there is no admissible cover

$$U = \bigcup_{i \in I} U_i \cup \bigcup_{j \in J} V_j,$$

with both summands nonempty, such that $U_i \cap V_j = \emptyset$ for $(i, j) \in I \times J$, as reflected by the fact that $\mathcal{O}(\mathbf{D}_K^1) = K\langle X \rangle$ is a domain.

Formalizing the above requires the notion of a *G-topology* on a topological space X , which is the data of a class of *admissible* open subsets¹⁰ and of *admissible* coverings of admissible open subsets satisfying some axioms. One has a natural notion of a sheaf with respect to a G-topology, which is a presheaf on the category of admissible opens which satisfies the



¹⁰ For \mathbf{D}_K^1 , we declare all open subsets admissible. The condition will however not be empty for \mathbf{D}_K^n with $n > 1$.

sheaf condition with respect to admissible coverings. Thus \mathcal{O} is a sheaf with respect to the admissible topology on \mathbf{D}_K^1 .

In Tate’s formalism, which we shall work out in the first part of the course, the basic geometric objects are *rigid-analytic varieties*. One uses as building blocks the *affinoid algebras*, which are quotients of the *Tate algebras*

$$K\langle X_1, \dots, X_r \rangle = \left\{ \sum_{n_1, \dots, n_r \geq 0} a_{n_1, \dots, n_r} X_1^{n_1} \dots X_r^{n_r} : a_{n_1, \dots, n_r} \rightarrow 0 \text{ as } n_1 + \dots + n_r \rightarrow \infty \right\}.$$

To an affinoid algebra $A = K\langle X_1, \dots, X_r \rangle / I$ one associates the *affinoid* $\text{Sp}A$. Its underlying topological space is the corresponding closed subset of

$$\mathbf{D}_K^r = \{(x_1, \dots, x_r) \in K^r : |x_i| \leq 1 \text{ for } i = 1, \dots, r\}$$

cut out by the ideal I . One equips it with a G-topology (the admissible topology), and a sheaf of rings \mathcal{O} , similarly to the case of \mathbf{D}_K^1 . A rigid-analytic variety is a topological space with a G-topology and a sheaf of rings with respect to that topology, which is locally (as a G-topologized space!) isomorphic to $\text{Sp}A$ for some affinoid algebra A .

1.3 Raynaud’s approach

The main drawbacks of Tate’s theory are

- the admissible topology is counterintuitive and complicated to work with,
- and the underlying spaces do not have enough points (e.g. there exist nonzero abelian sheaves for the admissible topology whose stalk at every point is zero),
- one is bound to work over a fixed field; for a non-algebraic extension of nonarchimedean fields K'/K (e.g. $\mathbf{C}_p/\mathbf{Q}_p$) there is no map $\mathbf{D}_{K'}^1 \rightarrow \mathbf{D}_K^1$,
- (why should there have to be a base field at all?)
- it is quite far from algebraic geometry (e.g. the opens are not defined by non-vanishing loci, but also be inequalities—not algebraic opens, but semi-algebraic opens).

There are several frameworks which address these issues in different ways, notably Huber’s theory of *adic spaces*, Berkovich’s theory of analytic spaces (usually called *Berkovich spaces*), and Raynaud’s approach via *formal schemes*, worked out by Bosch and Lütkebohmert and recently developed further by Fujiwara–Kato and Abbes. In the second half of this course, we will become acquainted with all of these, mostly focusing on Raynaud’s theory, as it is the closest to algebraic geometry.

THE STARTING POINT of Raynaud’s theory is the following isomorphism (where $t \in K$ is a pseudouniformizer)

$$K\langle X \rangle = \left(\varprojlim_m \mathcal{O}[X]/(t^m) \right) \left[\frac{1}{t} \right]. \tag{1.6}$$

We will prove this later, but you are welcome to try and check it yourself.

The isomorphism (1.6) expresses $K\langle X \rangle$ in terms of (0) the polynomial algebra $\mathcal{O}[X]$ through the algebraic operations of (1) t -adic completion, and (2) localization with respect to t . So, for example, if \mathcal{O} is a discrete valuation ring, we immediately see that $K\langle X \rangle$ is Noetherian, because (0) the polynomial algebra $\mathcal{O}[X]$ is Noetherian, (1) the completion of a Noetherian ring with respect to an ideal is Noetherian, and (2) the localization of a Noetherian ring is Noetherian. (Unfortunately, our \mathcal{O} will not always be Noetherian, so one needs to work harder.)

TO HAVE A GEOMETRIC PICTURE, we replace $\mathcal{O}[X]$ with its spectrum $X = \mathbf{A}_{\mathcal{O}}^1$. The projective system $\mathcal{O}/t^n \mathcal{O}[X]$ corresponds to a system of closed immersions

$$X_0 \hookrightarrow X_1 \hookrightarrow X_2 \hookrightarrow \cdots, \quad X_n = \mathbf{A}_{\mathcal{O}/t^{n+1}\mathcal{O}}^1.$$

Each of these immersions is defined a nilpotent ideal, and hence is a homeomorphism on the underlying spaces.

The above inductive system does not have a limit in the category of schemes. Instead, one can take its limit in the larger category of locally ringed spaces:

$$\mathfrak{X} = (|\mathfrak{X}|, \mathcal{O}_{\mathfrak{X}}) = \varinjlim_n X_n.$$

Since $|X_n| \hookrightarrow |X_{n+1}|$ are homeomorphisms, we can identify $|\mathfrak{X}|$ with $|X_0|$. Treating \mathcal{O}_{X_n} as a sheaf on $|X_0| = |\mathfrak{X}|$, we have

$$\mathcal{O}_{\mathfrak{X}} = \varprojlim_n \mathcal{O}_{X_n} = \varprojlim_n \mathcal{O}_X / (t^{n+1}).$$

The locally ringed space \mathfrak{X} is an example of a *formal scheme*, the *formal completion* of $X = \mathbf{A}_K^1$ with respect to the ideal $t \mathcal{O}_X$. In fact, in this context we could *define* formal schemes over \mathcal{O} as systems of closed immersions $X_0 \hookrightarrow X_1 \hookrightarrow \cdots$ between \mathcal{O} -schemes, with X_n defined by the ideal $t^{n+1} \mathcal{O}_{X_{n+1}}$.

The final step, inverting t , is the hardest: in Raynaud's approach, one wants to define a rigid-analytic variety over \mathcal{O} as the "generic fiber" of a formal scheme over \mathcal{O} . This is done purely formally by localizing the *category* of formal schemes over \mathcal{O} with respect to *admissible blow-ups*, i.e. blowups along an ideal containing a power of t . In the words of Fujiwara and Kato, *rigid geometry is the birational geometry of formal schemes*.

1.4 Why study rigid geometry?

The goal of the course is not only to introduce the basic definitions and facts surrounding rigid-analytic varieties—we will see some important applications of the theory as well. I will now try to give a short preview without spoilers.

Disclaimer: There are many possible answers to the question above. The following is heavily influenced by my own perspective and expertise as an algebraic geometer interested in the topology of algebraic varieties.

The broad answer is:

Rigid geometry allows us to use methods of topology and analysis in an otherwise purely algebraic context.

For an explicit example, consider a complex algebraic curve, say a smooth plane curve X in \mathbf{P}^2 of degree d . As one learns in the basic algebraic geometry course, this curve has genus

$$g = \frac{(d-1)(d-2)}{2}.$$

Over the complex numbers, the underlying manifold (the *complex analytification*) of X is an oriented surface with g many handles. Can we make sense of the last sentence algebraically? The question sounds crazy at first: to begin with, the underlying topological space of X (with the Zariski topology) does not see the genus at all, so how can we try to decompose it into handles?

Rigid geometry allows us to break varieties into pieces and perform surgery.

The answer is to *degenerate* the curve until it breaks and becomes easier to manage.¹¹ Thus, let ℓ_1, \dots, ℓ_d be generically chosen linear forms on \mathbf{P}^2 . If $\{f = 0\}$ is the homogeneous equation of our curve X , we consider the equation with an additional parameter t

$$X_t = \{tf + (1-t)\ell_1 \cdot \dots \cdot \ell_d = 0\} \subseteq \mathbf{P}_{k[t]}^2.$$

Thus $X_1 = X$, while X_0 is the union of d lines in \mathbf{P}^2 in general position.

The curve X_0 , while much easier to understand than X , is singular. Its topology differs from that of X . The idea, made possible by rigid geometry, is to study the smooth fibers X_t which “infinitesimally close” to X_0 . To make this precise, we first base change the above family to the field $K = k((t))$, obtaining a smooth algebraic curve X_K over K . Next, we turn it into a rigid-analytic variety $\mathcal{X} = (X_K)_{\text{an}}$, its *rigid analytification*. It is cut out by the same equation in a rigid-analytic version of \mathbf{P}_K^2 .

It turns out that \mathcal{X} is “close enough” to X_0 that there exists a natural morphism of topological spaces (the *specialization map*)

$$\text{sp}: |\mathcal{X}| \rightarrow |X_0|.$$

The preimage $U_i = \text{sp}^{-1}(L_i)$ of the line $L_i = \{\ell_i = 0\} \subseteq |X_0|$ happens to be an *open* rigid subvariety of \mathcal{X} which closely resembles a sphere with $d - 1$ discs removed (the discs are the preimages of the points $L_i \cap L_j$ for $j \neq i$ under sp). This gives a combinatorial decomposition of \mathcal{X} which one can use in place of the triangulation or handlebody decomposition on the complex analytification. For cubic curves (elliptic curves) one has the following picture:




Figure 1.1: Intuitive picture of the specialization map ($d = 3$, so $g = 1$).

¹¹ Can we study algebraic curves by putting them inside the Large Hadron Collider?

HERE ARE SOME EXAMPLES of serious applications of rigid geometry roughly along the above lines:

- Uniformization of curves and abelian varieties. (In fact, constructing a p -adic analytic analog of the expression of a complex elliptic curve as \mathbb{C} modulo a lattice was Tate's original motivation for defining rigid-analytic varieties. We will see Tate's uniformization later in the course.)
- The approach to SYZ mirror symmetry proposed by Kontsevich.
- Raynaud's solution to Abhyankar's conjecture (constructing finite étale covers of $\mathbb{A}_{\mathbb{F}_p}^1$ with given Galois group).
- Study of moduli of curves (often done using tropical methods, which is philosophically similar).
- Semistable reduction.

Other extremely important applications belong to p -adic Hodge theory.

2

Non-archimedean fields

In this chapter, we learn some fundamentals about the kind of base fields we will work with — fields complete with respect to a nontrivial non-archimedean norm. We start with basic facts about general valuation rings; the extra generality is not needed for Tate’s theory, but will prove useful later on.

In the appendix to this chapter, we review henselian local rings and Hensel’s lemma.

2.1 Valuation rings and valuations

Definition 2.1.1. A subring \mathcal{O} of a field K is a *valuation (sub)ring* of K if for every $x \in K^\times$, either $x \in \mathcal{O}$ or $x^{-1} \in \mathcal{O}$.

The above condition implies that $K = \text{Frac } \mathcal{O}$. This motivates the terminology: we will call a ring \mathcal{O} a *valuation ring* if \mathcal{O} is a domain and if it is a valuation ring of $K = \text{Frac } \mathcal{O}$.

Lemma 2.1.2. *Every valuation ring is a local ring.*

Proof. It suffices to check that the set of non-units is closed under addition. If $x, y \in \mathcal{O}$ are nonzero non-units, then either $xy^{-1} \in \mathcal{O}$, in which case $x + y = y(xy^{-1} + 1)$ is a non-unit because y is a non-unit, or $yx^{-1} \in \mathcal{O}$, and we swap x and y . \square

Lemma 2.1.3. *The relation*

$$x \leq y \quad \text{if} \quad yx^{-1} \in \mathcal{O} \tag{2.1}$$

*induces a linear order on $\Gamma = K^\times / \mathcal{O}^\times$, making Γ into a linearly ordered group.*¹

Proof. First, if $x' = ux$ and $y' = vx$ with $u, v \in R^\times$, then $x \leq y \iff x' \leq y'$, so that \leq induces a relation on $K^\times / \mathcal{O}^\times$. The fact that either $x \leq y$ or $y \leq x$ is the definition of a valuation ring. The rest is straightforward. \square

The quotient homomorphism

$$K^\times \rightarrow K^\times / \mathcal{O}^\times$$

is a “valuation” on the field K , as we shall now define. First, we introduce the following convention: for an ordered abelian group Γ (written additively), we shall write $\Gamma \cup \{\infty\}$ for the ordered commutative monoid

¹ An *ordered abelian group* is an abelian group Γ with an order relation \leq such that $a \leq b$ implies $a + c \leq b + c$. It is *linearly* or *totally ordered* if \leq is a linear order.

obtained by adding an element ∞ and declaring

$$\gamma \leq \infty \quad \text{and} \quad \gamma + \infty = \infty + \infty = \infty \quad (\gamma \in \Gamma).$$


Definition 2.1.4. A *valuation* on a field K is a group homomorphism

$$v: K^\times \rightarrow \Gamma$$

into a linearly ordered group Γ (written additively, so that $v(xy) = v(x) + v(y)$), which, when extended to a map of monoids $v: K \rightarrow \Gamma \cup \{\infty\}$ by $v(0) = \infty$, satisfies

$$v(x + y) \geq \min\{v(x), v(y)\}.$$

The *value group* of a valuation $v: K^\times \rightarrow \Gamma$ is the image $v(K^\times)$. Thus v trivially induces a surjective valuation $v': K^\times \rightarrow v(K^\times)$, and it is useful to identify v and v' . More generally, we will call two valuations $v_i: K^\times \rightarrow \Gamma_i$ ($i = 1, 2$) *equivalent* if there exists a third valuation $v: K^\times \rightarrow \Gamma$ and monotone homomorphisms $\varphi_i: \Gamma \rightarrow \Gamma_i$ ($i = 1, 2$) such that $v_i = \varphi_i \circ v$:



A valuation is *trivial* if it has trivial value group, i.e. $v(x) = 0$ for all $x \in K^\times$.

Proposition 2.1.5. Let K be a field.

(a) If $\mathcal{O} \subseteq K$ is a valuation ring and $\Gamma = K^\times / \mathcal{O}^\times$ is equipped with the linear order (2.1), then the projection map $v: K^\times \rightarrow \Gamma$ is a valuation on K .

(b) Conversely, if $v: K^\times \rightarrow \Gamma$ is a valuation, then

$$\mathcal{O} = \{x \in K \mid v(x) \geq 0\}$$

is a valuation ring of K , and its maximal ideal is $\mathfrak{m} = \{x \in K \mid v(x) > 0\}$.

(c) Constructions in (a) and (b) produce mutually inverse bijections

$$\{\text{valuation rings of } K\} \simeq \{\text{valuations on } K\} / \text{equivalence}.$$


Proof. (a) We check the property $v(x + y) \geq \min\{v(x), v(y)\}$, which resembles the proof that a valuation ring is local. Let $x, y \in K^\times$, and suppose $xy^{-1} \in \mathcal{O}$, then

$$v(x + y) = v(y(xy^{-1} + 1)) = v(y) + \underbrace{v(xy^{-1} + 1)}_{\geq 0 \text{ since } xy^{-1} + 1 \in \mathcal{O}} \geq v(y),$$

and similarly if $yx^{-1} \in \mathcal{O}$.

(b) Clearly for $x \in K$ either $x \in \mathcal{O}$ or $x^{-1} \in \mathcal{O}$ and \mathcal{O} is closed under multiplication. The fact that it is also closed under addition follows from $v(x + y) \geq \min\{v(x), v(y)\}$.

(c) Clearly, equivalent valuations define the same valuation ring. The only non-obvious assertion is that if $v_2: K^\times \rightarrow \Gamma_2 = K^\times / \mathcal{O}^\times$ is the valuation associated via (b) to the valuation ring \mathcal{O} associated to a valuation $v_1: K^\times \rightarrow \Gamma_1$ via (a), then v_1 and v_2 are equivalent. We let $\Gamma = \Gamma_2 = K^\times / \mathcal{O}^\times$, φ_1 the identity, and $\varphi_2: \Gamma = K^\times / \mathcal{O}^\times \rightarrow \Gamma_1$ the map induced by v_1 . \square



2.2 Valuations and norms

If the value group is a subgroup of \mathbf{R} , one can turn a valuation into a “norm.”

Definition 2.2.1. A valuation of height one² is a valuation $v: K^\times \rightarrow \mathbf{R}$.

Note that two valuations of height one $v_i: K^\times \rightarrow \mathbf{R}$ ($i = 1, 2$) are equivalent if and only if $v_2(x) = c v_1(x)$ for some positive real c .³

Definition 2.2.2. A nonarchimedean norm on a field K is a map

$$|\cdot|: K \rightarrow [0, \infty)$$

such that

- i. $|xy| = |x| \cdot |y|$,
- ii. $|x| = 0$ if and only if $x = 0$,
- iii. $|x + y| \leq \max\{|x|, |y|\}$.

Proposition 2.2.3. Let K be a field.

(a) If $v: K \rightarrow \mathbf{R}$ is valuation of height one, then⁴

$$|x| = \exp(-v(x))$$

(where $\exp(-\infty) = 0$) defines a nonarchimedean norm on K .

(b) Conversely, if $|\cdot|$ is a norm on K , then

$$v(x) = -\log |x|$$

(where $\log 0 = -\infty$) defines a valuation of height one. The corresponding valuation ring is the “closed ball” $\mathcal{O} = \{x \mid |x| \leq 1\}$.

(c) The constructions in (a) and (b) produce mutually inverse bijections

$$\{\text{height one valuations on } K\} \simeq \{\text{nonarchimedean norms on } K\}.$$

Proof. Clear. \square

Proposition 2.2.4. Let $|\cdot|$ be a nonarchimedean norm on a field K . Then

$$d(x, y) = |x - y|$$

defines a metric on K , making K into a topological field. This metric and the induced topology have the following properties:

- (a) Every triangle is isosceles, every point of an open ball is its center, and every two (open or closed) balls are either disjoint or one contains the other,

² This terminology is slightly nonstandard: what is usually meant by a valuation of height one is a nontrivial valuation whose value group embeds in \mathbf{R} .

More generally, the height (or rank) of a valuation is the order type of the set of all convex subgroups of the value group, (linearly) ordered by inclusion, where a subgroup $A \subseteq \Gamma$ is convex if $a \leq x \leq b$ and $a, b \in A$ implies $x \in A$.

As it turns out, and is easy to show, this is just the Krull dimension of the corresponding valuation ring \mathcal{O} .

³ Exercise 3 on Problem Set 1.

⁴ The base e of the exponential is of course an arbitrary choice. Sometimes there exists a more natural one. For example, if K is p -adic, i.e. $|p| < 1$ for a prime p , then one usually considers the norm

$$|x| = p^{-v(x)}.$$

(b) The open ball $\{|x - a| < \rho\}$, the closed ball $\{|x - a| \leq \rho\}$, and the sphere $\{|x - a| = \rho\}$ are both open and closed for $\rho > 0$. In particular, the valuation ring $\mathcal{O} = \{|x| \leq 1\} \subseteq K$ is an open subring.

(c) The topological space K is totally disconnected,

(d) Suppose that K is complete (every Cauchy sequence converges). A series $\sum_{n=0}^{\infty} a_n$ with $a_n \in K$ converges if and only if $\lim a_n = 0$.

Proof. Continuity of addition, multiplication, and inverse is clear and left to the reader.

(a) The key observation is that if $|x| > |y|$, then $|x - y| = \max\{|x|, |y|\} = |x|$. Indeed, we have

$$|x| = |y + (x - y)| \leq \max\{|y|, |x - y|\} \leq \max\{|y|, |x|, |y|\} = |x|,$$

so the inequalities are equalities, implying $|x - y| = |x|$. Similarly, if $|y| > |x|$ then $|x - y| = |y|$, thus in general two of the numbers $|x|, |y|, |x - y|$ have to be equal.

If a triangle has vertices a, b, c , apply the above to $x = c - a, y = c - b$ to see that it is isosceles, with two longest sides being equal.

Now consider an open ball $B(a, \rho) = \{|x - a| < \rho\}$ and let $b \in B$, i.e. $|b - a| < \rho$. If $c \in K$, then consider the triangle with vertices a, b, c . The above observation shows that $|c - a| \geq \rho$ if and only if $|c - b| \geq \rho$, showing $B(a, \rho) = B(b, \rho)$.

If two open balls B and B' intersect at a point b , then taking b as the center of both balls shows that one is contained in the other.

(b) The open ball is of course open, and the closed ball is the union of the open ball and the sphere. It suffices to treat the sphere $S = \{|x| = \rho\}$ (centered at zero for simplicity). Let $a \in S$; we claim that the open ball $\{|x - a| < \rho\}$ is contained in S . Indeed, if $|x - a| < \rho$ then $|x| = |a + (x - a)|$ and since $|x - a| < \rho = |a|$, we have $|x| = |a| = \rho$, so $x \in S$.


(c) Let $S \subseteq K$ be a subset and let $a, b \in S$ be two distinct points, $\rho = |a - b| > 0$. Then

$$S = (S \cap \{|x - a| < \rho/2\}) \cup (S \cap \{|x - a| \geq \rho/2\})$$

expresses S as a sum of two disjoint and non-empty open subsets. Thus S cannot be connected if it has more than one point.

(d) Clearly if $\sum a_n$ converges then $\lim a_n = 0$. Conversely, suppose $\lim a_n = 0$; we check that $b_n = a_1 + \dots + a_n$ forms a Cauchy sequence. Let $\varepsilon > 0$, and let N be such that $|a_n| < \varepsilon$ for $n \geq N$. Then for $m > n > N$

$$|b_m - b_n| = |a_{n+1} + \dots + a_m| < \max\{|a_{n+1}|, \dots, |a_m|\} < \varepsilon. \quad \square$$



2.3 Geometric examples of valuations

Long long time ago, before schemes were invented by Grothendieck, varieties were studied (or even defined) using valuations on their function fields. E.g. Zariski's proof of resolution of singularities on surfaces heavily relied on the classification of valuations on their function fields. We will see some of these below.

This section is a bit of a digression, but will become important later in the course.

Example 2.3.1. Let R be a Dedekind domain with field of fractions K , and let $\mathfrak{m} \subseteq R$ be a maximal ideal. Standard examples:

- $R = \Gamma(X, \mathcal{O}_X)$ for X a smooth affine algebraic curve, with \mathfrak{m} corresponding to a closed point $x \in X$,
- $R = \mathcal{O}_K$ the ring of integers in a number field K , e.g. $R = \mathbf{Z}[i]$.

The local ring $\mathcal{O} = R_{\mathfrak{m}}$ is a discrete valuation subring of K . The corresponding valuation on K is $v(x) = \max\{k : x \in \mathfrak{m}^k\}$. Every valuation on K which is trivial on k is equivalent to exactly one of these.⁵

⁵ Sound familiar? [7, Chapter I 6]

The remaining examples deal valuations on function fields of surfaces over a base field k , where the situation is much more complicated, essentially due to the existence of non-trivial blowups.⁶ We only consider valuations whose restriction to k is trivial.

⁶ See [7, Exercise II 4.12].

Example 2.3.2 (Divisorial valuation). Let X be a normal surface with field of rational functions K and let $D \subseteq S$ be a prime divisor. Then [7, II 6] D defines a function “order of zero along D ”

$$v_D : K = k(S) \rightarrow \mathbf{Z} \cup \{\infty\}$$

which is a valuation. The corresponding valuation ring is $\mathcal{O}_{X,\xi}$ where ξ is the generic point of D . Its residue field is $k(D)$, the function field of D .

Example 2.3.3 (Valuation of height two). In the situation of Example 2.3.2, let $p \in D$ be a closed point at which D is regular. Then x defines a valuation v_p on $k(D)$ as in Example 2.3.1. We can combine the valuations v_D on $K = k(S)$ and v_p on $k(D)$ into a height two valuation

$$v_{D,p} : K \rightarrow \mathbf{Z}_{\text{lex}}^2 \cup \{\infty\},$$

where $\mathbf{Z}_{\text{lex}}^2$ is \mathbf{Z}^2 with the lexicographic order $((x, y) \geq (x', y')$ if $x > x'$ or $x = x'$ and $y \geq y')$. To define $v_{D,p}$, we pick a uniformizer (generator of the maximal ideal) $\pi \in \mathcal{O}_{X,\xi} = \mathcal{O}_{v_D}$ without zero or pole at p and set

$$v_{D,p}(f) = (v_D(f), v_p(g)), \quad g = (\pi^{-v_D(f)} f)|_{\xi},$$

where the restriction makes sense because $v_D(g) = 1$, so $g \in \mathcal{O}_{v_D}$.

The valuation ring $\mathcal{O}_{v_{D,p}}$ consists of rational functions with no pole along D and whose restriction to D has no pole at p . It has three prime ideals, is of Krull dimension two, and is non-Noetherian. Its residue field is k . See Figure 2.1 for the monoid of monomials in $\mathcal{O}_{v_{D,p}}$ for $S = \mathbf{A}^2$.

Example 2.3.4 (Valuations from formal curve germs). Let again S be a normal surface with function field K , and let

$$\gamma : \text{Spec } k[[t]] \rightarrow S$$

be a morphism of schemes (a “formal curve germ”). We say that γ is *nonalgebraic* if its image is not contained in a proper closed subscheme of S , equivalently if γ maps the generic point $\text{Spec } k((t))$ of $\text{Spec } k[[t]]$ to the generic point $\eta = \text{Spec } K$ of S .⁷ The composition of γ^* with the standard

⁷ There is plenty of nonalgebraic curve germs on an algebraic surface. For example, consider $S = \text{Spec } \mathbf{C}[x, y]$ the affine plane and γ defined by

$$\gamma^*(x) = t, \quad \gamma^*(y) = \exp t = \sum_{n \geq 0} \frac{t^n}{n!}.$$




Figure 2.1: In Example 2.3.3, consider $S = \mathbf{A}^2$ with coordinates x, y , the divisor $D = \{x = 0\} \subseteq S$, and the point $p = \{y = 0\} \subseteq D$. The figure shows the monoid consisting of all $(m, n) \in \mathbf{Z}^2$ for which $v(x^m y^n) \geq 0$. Can you see why this monoid is not finitely generated? This is related to the fact that the valuation ring is non-Noetherian.

valuation on $k((t))$ gives a height one valuation

$$v_\gamma : K \rightarrow k((t)) \rightarrow \mathbf{Z} \cup \{\infty\}$$

with residue field k .

Example 2.3.5 (Height one valuation with dense value group). Suppose that $K = k(x, y)$. Let λ be an irrational real number. Define the weight function on monomials in x and y by

$$\text{weight}_\lambda(x^m y^n) = m + \lambda n \in \mathbf{R}.$$

Define the valuation $v_\lambda : K \rightarrow \mathbf{R} \cup \{\infty\}$ by first defining it on polynomials:

$$v_\lambda \left(\sum_{m,n \geq 0} a_{mn} x^m y^n \right) = \min \{ \text{weight}_\lambda(x^m y^n) : a_{mn} \neq 0 \}$$

and extending to $k(x, y)$ by $v_\lambda(f/g) = v_\lambda(f) - v_\lambda(g)$. This gives a valuation on K which has height one but whose value group $\mathbf{Z} \oplus \lambda \mathbf{Z} \simeq \mathbf{Z}^2$ is dense in \mathbf{R} . See Figure 2.2 for the monoid of monomials in the valuation ring.

Remark 2.3.6. The valuation v_λ in Example 2.3.5 can be thought of as the valuation of the type considered in Example 2.3.4 induced by the “formal curve germ”

$$t \mapsto (t, t^\lambda).$$

In fact, for $\lambda' = a/b$ rational with $(a, b) = 1$, we can define the curve germ

$$\gamma_{a,b} : \text{Spec } \mathbf{C}[[t]] \rightarrow \mathbf{A}_{x,y}^2, \quad \gamma_{a,b}^*(x) = t^b, \quad \gamma_{a,b}^*(y) = t^a.$$

Let $v_{a,b} = \frac{1}{b} v_{\gamma_{a,b}}$ where $v_{\gamma_{a,b}}$ is the valuation associated to the curve germ as in Example 2.3.4. If $a_n/b_n \rightarrow \lambda$, then the corresponding valuations v_{a_n, b_n} converge pointwise to v_λ .

2.4 Nonarchimedean fields

Definition 2.4.1. A *nonarchimedean field*⁸ is a field K equipped with a nontrivial nonarchimedean norm $|\cdot|$ with respect to which it is complete.

⁸ For many authors, “nonarchimedean field” is simply a field with a nonarchimedean norm.




Figure 2.2: The monoid of all $(m, n) \in \mathbf{Z}^2$ for which $v(x^m y^n) \geq 0$ (Example 2.3.5). The boundary of the gray area is the line with slope $-1/\lambda$

$$x + \lambda y = 0.$$

Since $\lambda \notin \mathbf{Q}$, this line contains no nonzero lattice points.

Proposition 2.4.2. *Let K be a field endowed with a nontrivial nonarchimedean norm $|\cdot|$. The ring operations on K extend uniquely to the completion \widehat{K} of K with respect to $d(x, y) = |x - y|$, making \widehat{K} into a nonarchimedean field.*

Definition 2.4.3. Let K be a field endowed with a nonarchimedean norm $|\cdot|$. A *pseudouniformizer* is an element $t \in K$ with $0 < |t| < 1$.⁹

Thus $|\cdot|$ is nontrivial if and only if K admits a pseudouniformizer.

Proposition 2.4.4. *Let K be a field endowed with a nontrivial nonarchimedean norm $|\cdot|$, and let $t \in K$ be a pseudouniformizer. Let $\mathcal{O} = \{x \in K \mid |x| \leq 1\}$ be the valuation ring. Then K is complete (i.e. K is a nonarchimedean field) if and only if \mathcal{O} is t -adically complete and separated, i.e. if the natural map*

$$\pi: \mathcal{O} \rightarrow \varprojlim_n \mathcal{O}/t^n \mathcal{O}$$

is an isomorphism. In this case, the map π is a homeomorphism, where the target is endowed with the inverse limit topology where each $\mathcal{O}/t^n \mathcal{O}$ is given the discrete topology.

Proof. Set $\rho = |t|$; we have $0 < \rho < 1$. First, we note that

$$t^n \mathcal{O} = \{x \in K : |x| \leq \rho^n\}.$$

The kernel of π is $\bigcap_{n \geq 0} t^n \mathcal{O} = \{|x| \leq 0\} = \{0\}$. Thus π is always injective.

An element \bar{f} of the inverse limit is a compatible system $(\bar{f}_n \in \mathcal{O}/t^n \mathcal{O})$. Let $f_n \in \mathcal{O}$ be elements mapping to $\bar{f}_n \in \mathcal{O}/t^n \mathcal{O}$. We claim that (f_n) is a Cauchy sequence. Indeed, we have $f_n - f_m \in t^n \mathcal{O}$ for $m > n$, so $|f_n - f_m| \leq \rho^n$ for $m > n$. Thus if K is complete, then (f_n) has a limit $f \in \mathcal{O}$. Now for every n , we have

$$|f - f_n| = |f_n - f_m| \leq \rho^n \quad \text{for } m \gg 0,$$

which shows that $f - f_n \in t^n \mathcal{O}$. Thus $\pi(f) = \bar{f}$, i.e. π is surjective if K is complete.

⁹ In other words, t is a *topologically nilpotent unit*, where topologically nilpotent means that $|t^n| \rightarrow 0$.

Warning: if K is not discretely valued, then \mathcal{O} will not be a complete local ring! In that case, the maximal ideal of \mathcal{O} satisfies $\mathfrak{m}^2 = \mathfrak{m}$, and hence $\mathcal{O}/\mathfrak{m}^n = k$ for all n , so that $\widehat{\mathcal{O}} \simeq k$. This is why we need to work with pseudouniformizers.

Conversely, suppose that π is surjective. We will show that \mathcal{O} is complete with respect to $|\cdot|$ (this easily implies that K is complete). Let $(f_n) \in \mathcal{O}$ be a Cauchy sequence. For every m , the images of f_n in $\mathcal{O}/t^m \mathcal{O}$ have to stabilize for $n \gg 0$. Let $\bar{f}_m \in \mathcal{O}/t^m \mathcal{O}$ be the stable value (i.e. $\bar{f}_m = \lim_n (f_n \bmod t^m)$ for the discrete topology on $\mathcal{O}/t^m \mathcal{O}$). It is easy to see that $f = (\bar{f}_m)$ is an element of the inverse limit of $\mathcal{O}/t^n \mathcal{O}$. Let $f \in \mathcal{O}$ be an element with $\pi(f) = \bar{f}$, then $f = \lim f_n$.

The claim about the topologies follows from the fact that $t^n \mathcal{O} = \{|x| \leq \rho^n\}$ is a basis of neighborhoods of zero in \mathcal{O} . □

2.5 Extensions of nonarchimedean fields

The treatment here follows [4, Appendix A] and [8, II §4 and §6].

Theorem 2.5.1. *Let K be a nonarchimedean field and let L/K be a finite extension. Then there exists a unique norm $|\cdot|$ on L extending K . The field L endowed with this norm is a nonarchimedean field.*

For $f = \sum_{i=0}^n a_i x^i \in K[X]$, we define its *Newton polygon* $\text{NP}(f)$ as the lower convex envelope of the set $\{(0, v(a_0)), \dots, (n, v(a_n))\}$ in \mathbf{R}^2 . Its basic property is that $\text{NP}(fg) = \text{NP}(f) + \text{NP}(g)$ (Minkowski sum, i.e. sort the segments of both polygons by slope and concatenate). In particular, if f is reducible, then $\text{NP}(f)$ contains a point of the form (m, γ) with $0 < m < \deg f$ an integer and γ an element of the value group. One form of Hensel's lemma¹⁰ states a partial converse:

Lemma 2.5.2 (Irreducibility and Newton polygons). *Let $f \in K[X]$ be a nonzero polynomial with $f(0) \neq 0$. Then f is irreducible if $\text{NP}(f)$ is a single segment without interior points of the form (m, γ) with $m \in \mathbf{Z}$ and $\gamma \in v(K^\times)$. Conversely:*

- (a) (Weak form) *If $\text{NP}(f)$ has segments both of negative and of non-negative slope, then f is reducible.*
- (b) (Strong form) *If f is irreducible, then $\text{NP}(f)$ is a single segment.*

We shall prove the weak form now. It will be sufficient for the proof of Theorem 2.5.1, which in turn will be used to prove the strong form.

Proof (of the weak form). The first assertion has already been explained in the discussion preceding the statement of the lemma. To show (a), let (m, γ) be a vertex of $\text{NP}(f)$ with smallest γ , and with smallest m among those. Then $0 < m < \deg f$, otherwise all slopes of $\text{NP}(f)$ have the same sign (see Figure 2.5). Replacing f with $a_m^{-1} f$, we may assume that $\gamma = 0$, and consequently $f \in \mathcal{O}[X]$. The image \bar{f} of f in $k[X]$ decomposes as

$$\bar{f} = X^m b(X) \quad \text{with} \quad b(0) \neq 0.$$

By Hensel's lemma (Proposition 2.A.5) using the formulation as in Proposition 2.A.1(b), the above factorization lifts to a factorization $f = \tilde{g} \tilde{h}$ with $\deg \tilde{g} = m$. Therefore f is reducible. □

Proposition 2.5.3. *In the situation of Theorem 2.5.1, let $\mathcal{O} = \{|x| \leq 1\}$ be the valuation ring of K . An element $x \in L$ is integral over \mathcal{O} if and only if $\text{Nm}_{L/K}(x) \in \mathcal{O}$.*




Figure 2.3: Newton polygon of the polynomial

$$1 + \pi^{-1}X - \pi^{-1}X^2 + \pi X^3 + \pi^2 X^5$$

¹⁰ In the appendix to this lecture, we shall discuss different formulations of Hensel's lemma.




Figure 2.4: Proof of Lemma 2.5.2(a)

Proof. Let $f \in K[X]$ be the minimal polynomial of x . Since f is irreducible, by Lemma 2.5.2 its Newton polygon has to be the line segment with endpoints $(\deg f, 0)$ and $(0, c)$ where $c = v(a_0)$ is the valuation of the constant term of f (Figure 2.5). But $c = (-1)^n \text{Nm}_{L/K}(x)$, so if $\text{Nm}_{L/K}(x) \in \mathcal{O}_K$ then $\text{NP}(f)$ lies entirely above the line $y = 0$, which implies that $f \in \mathcal{O}[X]$, so that x is integral over \mathcal{O} .

Conversely, if x is integral, then in fact its minimal polynomial f belongs to $\mathcal{O}[X]$; in particular, $\text{Nm}_{L/K}(x) = (-1)^{\deg f} f(0) \in \mathcal{O}$. To see this, let $g \in \mathcal{O}[X]$ be monic with $g(x) = 0$. We have $g = fh$ for some (also monic) $h \in K[X]$. Then $\text{NP}(g) = \text{NP}(f) + \text{NP}(h)$ lies above the line $y = 0$ and ends on it (because it is monic), and hence all of its slopes are non-positive. However, $\text{NP}(f)$ is a single segment (connecting $(0, c)$ and $(\deg f, 0)$), and its slope is one of the slopes of $\text{NP}(g)$ and hence is non-positive. Thus $c \geq 0$, i.e. $f \in \mathcal{O}[X]$. \square




Figure 2.5: Newton polygon of an irreducible monic polynomial f (Proof of Proposition 2.5.3)

Proof of Theorem 2.5.1. Let $\mathcal{O} = \{|x| \leq 1\} \subseteq K$ be the valuation ring of K and let $\mathcal{O}' \subseteq L$ be the integral closure of \mathcal{O} inside L . By Proposition 2.5.3, $x \in \mathcal{O}'$ if and only if $|\text{Nm}_{L/K}(x)| \leq 1$. Since the norm is multiplicative, this shows that \mathcal{O}' is a valuation ring of L . Moreover, $\mathcal{O}' \cap K = \mathcal{O}$ because \mathcal{O} is integrally closed.¹¹

Define $|x| = |\text{Nm}_{L/K}(x)|^{1/d}$ for $x \in L$, where $d = [L : K]$. This restricts to the norm on K , is multiplicative, and $|x| \neq 0$ for $x \neq 0$. To show $|x + y| \leq \max\{|x|, |y|\}$, we use the fact that $\{|x| \leq 1\} = \mathcal{O}'$ is a valuation ring.

If $|\cdot|'$ is some other norm extending $|\cdot|$ to L , then since the corresponding valuation ring $\{|x'| \leq 1\}$ is integrally closed, it contains \mathcal{O}' . This implies that $|\cdot| \leq |\cdot|'$, and by Exercise 3 from Problem Set 1, we have $|\cdot|' = |\cdot|^c$ for some constant c . But $c = 1$ since the two agree on K . \square

Theorem 2.5.4 (Krasner). *Let K be a nonarchimedean field, and let \bar{K} be an algebraic closure of K , which we endow with the unique extension of $|\cdot|$. Let $\widehat{\bar{K}}$ denote the completion of \bar{K} with respect to this norm. Then $\widehat{\bar{K}} = \bar{K}^\wedge$ is algebraically closed.*

Proof. Let L be a finite extension of $\widehat{\bar{K}}$. By Theorem 2.5.1, there exists a unique norm on L extending the norm on $\widehat{\bar{K}}$ and L is complete with respect to that norm. To show $L = \widehat{\bar{K}}$, it therefore suffices to prove that $\widehat{\bar{K}}$ is dense in L .

Let $x \in L$ and let $1 > \rho > 0$. We shall find a $y \in \widehat{\bar{K}}$ with $|x - y| < \rho$. Without loss of generality, we may assume that $|x| \leq 1$. Let $f = \sum_{i=0}^n a_i X^i \in \widehat{\bar{K}}[X]$ be its minimal polynomial (with $a_n = 1$). Since \bar{K} is dense in $\widehat{\bar{K}}$, we can find $b_i \in \bar{K}$ ($i = 0, \dots, n$) with $|a_i - b_i| < \rho$ (and again $b_n = 1$). This implies that

$$|g(x)| = |g(x) - f(x)| = \left| \sum_{i=0}^n (a_i - b_i)x^i \right| < \rho.$$

Now, the polynomial $g = \sum_{i=0}^n b_i X^i$ splits completely in \bar{K} :

$$g = \prod_{i=1}^n (X - \gamma_i), \quad \gamma_1, \dots, \gamma_n \in \bar{K}.$$

¹¹ Easy exercise: show that every valuation ring is integrally closed.

Evaluating at x and taking absolute value, we obtain

$$\rho > |g(x)| = \prod_{i=1}^n |x - y_i|.$$

Therefore one of the factors is less than ρ . \square

2.6 Slopes of the Newton polygon

We can now prove the promised strong form of Lemma 2.5.2. It will not be used later in the course.

Lemma 2.6.1. *If $f \in K[X]$ is irreducible, then all roots of f in \overline{K} have the same norm.*

Proof. Let L/K be the splitting field of f and let $G = \text{Gal}(L/K)$. Thus G acts transitively on the roots of f in L . Since the norm $|\cdot|$ on L extending the norm on K is unique, the group G acts on L by isometries. In particular, for any two roots α, β of f in L we can find $g \in G$ with $\beta = g(\alpha)$, and then

$$|\alpha| = |g(\alpha)| = |\beta|. \quad \square$$

For a real number λ and $f \in K[X]$, we define the *slope multiplicity* $\mu(\lambda, f)$ of λ in $\text{NP}(f)$ as the length of the projection on the x -axis of the segment in $\text{NP}(f)$ with slope λ (zero if it does not exist), see Figure 2.6. Additivity of Newton polygons means precisely that

$$\mu(\lambda, fg) = \mu(\lambda, f) + \mu(\lambda, g) \quad \text{for every } \lambda \in \mathbf{R}.$$

Lemma 2.6.2. *For $f \in K[X]$ and $r > 0$, we have*

$$\#\{\alpha \in \overline{K} : f(\alpha) = 0 \text{ and } |\alpha| = r\} = \mu(\log r, f).$$

Proof. By additivity of both sides of the asserted equality, we may assume that f is irreducible, in which case all roots of f have the same absolute value ρ by Lemma 2.6.1. We may also assume that f is monic and $\rho \neq 0$, and write

$$f = \sum_{i=0}^n a_{n-i} X^i = \prod_{j=1}^n (X - \alpha_j), \quad |\alpha_j| = \rho.$$

Therefore for $0 < i \leq n$ we have

$$a_i = (-1)^i \sum_{0 \leq j_1 < \dots < j_i \leq n} \alpha_{j_1} \cdots \alpha_{j_i},$$

and taking absolute values we obtain

$$|a_i| \leq \rho^i \quad \text{and} \quad |a_n| = |\alpha_1 \cdots \alpha_n| = \rho^n.$$

It follows that $\text{NP}(f)$ is the segment connecting the points $(0, v(a_n)) = (0, -n \log \rho)$ and $(n, 0)$. This implies the asserted equality for $\rho = r$, with both sides equal to $n = \deg f$. Therefore for $r \neq \rho$ both sides are zero, and hence the assertion is true for every $r > 0$. \square

Proof of the strong form of Lemma 2.5.2. Let $f \in K[X]$ be irreducible. By Lemma 2.6.1, all roots of f have the same absolute value. By Lemma 2.6.2, the Newton polygon $\text{NP}(f)$ has a single slope, i.e. it is a segment. \square




Figure 2.6: Slope multiplicity

2.A Henselian rings

Hensel’s lemma played an important in the proof of Theorem 2.5.1. The first goal of this section is to elucidate its role by introducing the notion of a *henselian local ring*. Roughly speaking, it is a local ring in which the assertion of Hensel’s lemma holds. There are however many equivalent characterizations of this class of local rings, reviewed in Proposition 2.A.1 below, and the reader familiar with the étale topology will surely appreciate the topological flavor of some of them. The second goal is to prove Hensel’s lemma in its general form: *a local ring complete with respect to a \mathfrak{m} -primary ideal is henselian*.

Our treatment follows the Stacks Project [10, Tag 04GE].

Proposition 2.A.1. *Let A be a local ring with maximal ideal \mathfrak{m} . We set $k = A/\mathfrak{m}$, $x = \text{Spec}k$, $X = \text{Spec}A$, $i : x \rightarrow X$ the inclusion. The following conditions are equivalent:*

- (a) *If $f \in A[T]$ is monic and $t_0 \in k$ is a root of $\bar{f} = f \bmod \mathfrak{m} \in k[T]$ such that $\bar{f}'(t_0) \neq 0$, then there exists a unique root $t \in A$ of f such that $t \bmod \mathfrak{m} = t_0$.*
- (b) *If $f \in A[T]$ is monic and $\bar{f} = gh$ is a factorization of $\bar{f} = f \bmod \mathfrak{m} \in k[T]$ with $g, h \in k[T]$ coprime, then there exists a factorization $f = \tilde{g}\tilde{h}$ with $\tilde{g}, \tilde{h} \in A[T]$ such that $\tilde{g} \bmod \mathfrak{m} = g$, $\tilde{h} \bmod \mathfrak{m} = h$, and $\deg \tilde{g} = \deg g$.*
- (c) *Every finite A -algebra is a product of local rings.*
- (d) *For every étale A -algebra B and every prime $\mathfrak{p} \subseteq B$ lying over \mathfrak{m} and such that $k(\mathfrak{p}) = k$, there exists a section $s : B \rightarrow A$ of $A \rightarrow B$ with $\mathfrak{p} = s^{-1}(\mathfrak{m})$.*
- (e) *For every étale morphism $f : U \rightarrow X$ and every lifting $\tilde{i} : x \rightarrow U$ of i (i.e. $i = f \circ \tilde{i}$) there exists a unique section $s : X \rightarrow U$ such that $s \circ i = \tilde{i}$.¹²*

Proof. Maybe I’ll write something here later. □

Definition 2.A.2. (a) A local ring A is *henselian* if the equivalent conditions of Proposition 2.A.1 hold.

(b) A local ring A is *strictly henselian* if it is henselian and its residue field k is separably closed.¹³

(c) A valued field (K, ν) is *henselian* if the valuation ring $\mathcal{O} = \{x \mid \nu(x) \geq 0\}$ is henselian.

Remark 2.A.3. Condition (d) of Proposition 2.A.1 allows one to construct the *henselization* of a local ring A as the direct limit


$$A^h = \varinjlim_{(B,s) \in \mathcal{C}_A} B$$

where \mathcal{C}_A is the category of pairs (B, s) with B an étale A -algebra and $s : B \rightarrow k$ a homomorphism extending $A \rightarrow k$. (This category is filtering and essentially small.)

The ultimate reference is Raynaud’s book *Anneaux locaux henséliens*.

[10, Tag 04GG]

¹² Useful to picture this condition as a lifting problem:



¹³ Equivalently: every étale cover of $\text{Spec}A$ admits a section.

Universal property: $A \rightarrow A^b$ is a local homomorphism into a henselian local ring which is initial among such (in the category of local rings and local homomorphisms).

Similarly, given a separable closure k^{sep} of k , we can construct the *strict henselization* A^{sh} by considering the category of étale A -algebras endowed with a homomorphism to k^{sep} extending $A \rightarrow k^{\text{sep}}$. (Using the algebraic closure \bar{k} instead of k^{sep} gives the same result.)

Remark 2.A.4. The strict henselization of a local ring is the local ring for the étale topology. To make this precise, we reformulate everything in terms of geometry. Recall that a *geometric point* of a scheme X is a map $\bar{x} \rightarrow X$ with $\bar{x} = \text{Spec } k(\bar{x})$ for some separably closed field $k(\bar{x})$. (Again, one can use algebraically closed fields instead.) An *étale neighborhood* of a geometric point \bar{x} of X is an étale morphism $U \rightarrow X$ endowed with a lifting $\bar{x} \rightarrow U$ of $\bar{x} \rightarrow X$. Étale neighborhoods of \bar{x} in X form a cofiltering category $N(X, \bar{x})$, and the colimit

$$\mathcal{O}_{X, \bar{x}} = \varinjlim_{U \in N(X, \bar{x})} \Gamma(U, \mathcal{O}_U)$$

is isomorphic to the strict henselization $\mathcal{O}_{X, x}^{\text{sh}}$ of $\mathcal{O}_{X, x}$ where x is the image of \bar{x} in X (and where we use the separable closure of $k(x)$ in $k(\bar{x})$ as $k(x)^{\text{sep}}$).¹⁴

Proposition 2.A.5 (Hensel’s lemma). *Every local ring A which is J -adically complete and separated for an \mathfrak{m} -primary¹⁵ ideal $J \subseteq A$ is henselian. In particular, every complete local ring is henselian.*

For fans of the étale topology, we give a geometric proof:

Proof. We prove condition (e). Let $X = \text{Spec } A$ and $x = \text{Spec } k$ as before, and let

$$\begin{array}{ccc} & U & \\ \tilde{i} \nearrow & \downarrow f & \\ x & \xrightarrow{i} & X \end{array}$$

be an étale neighborhood of $x \rightarrow X$. Set $X_n = \text{Spec } A/J^{n+1}$ for $n \geq 0$. First, consider the diagram

$$\begin{array}{ccc} x & \xrightarrow{\tilde{i}} & U \\ \downarrow & \nearrow s_0 & \downarrow f \\ X_0 & \longrightarrow & X \end{array}$$

Since $x \rightarrow X_0$ is an immersion defined by the nil ideal¹⁶ $\mathfrak{m}/J \subseteq A/J$, by the infinitesimal criterion for étaleness¹⁷ there exists a unique diagonal arrow s_0 making the square commute.

Starting from s_0 , we shall successively build maps $s_n: X_n \rightarrow U$ lifting $X_n \rightarrow X$ along f . It suffices to apply the infinitesimal criterion to the squares

$$\begin{array}{ccc} X_n & \xrightarrow{s_n} & U \\ \downarrow & \nearrow s_{n+1} & \downarrow f \\ X_{n+1} & \longrightarrow & X \end{array}$$

¹⁴ Similarly, the henselization is related in the same way to local rings for the Nisnevich topology.

¹⁵ This means that for $x \in \mathfrak{m}$ we have $x^N \in J$ for $N \gg 0$ depending on x .

¹⁶ An ideal in a commutative ring is *nil* (locally nilpotent in [10]) if it consists of nilpotent elements.

¹⁷ **Infinitesimal criterion for étale maps:** A morphism $f: X \rightarrow Y$ locally of finite presentation is étale if and only if for every ring A and nil ideal $I \subseteq A$ (equivalently, every square zero ideal), and every commutative square of solid arrows

$$\begin{array}{ccc} \text{Spec } A/I & \longrightarrow & X \\ \downarrow & \nearrow & \downarrow f \\ \text{Spec } A & \longrightarrow & Y \end{array}$$

there exists a unique dotted arrow making the diagram commute.

Since A is J -adically complete, in the limit, the maps give the desired section $s: X \rightarrow U$.¹⁸ □

¹⁸ If you are confused with the last step, set $U = \text{Spec } B$ and temporarily revert to commutative algebra.

Remark 2.A.6. The most common proof uses condition (a) of Proposition 2.A.1, and uses “Newton’s method” to iteratively construct the desired root t using explicit induction steps. Proofs in [4, Appendix A] and [8] use condition (b), which gives a more direct approach to proving Theorem 2.5.1, but makes for a messier and less illuminating argument.

Corollary 2.A.7. *Every nonarchimedean field is henselian.*

Proof. Let K be a nonarchimedean field, let $\mathcal{O} \subseteq K$ be its valuation ring, and let $t \in \mathcal{O}$ be a pseudouniformizer. Apply Proposition 2.A.5 with $A = \mathcal{O}$ and $J = (t)$. □

Lemma 2.A.8. *The following are equivalent for a field K endowed with a height one valuation ν .*

(a) K is henselian.

(b) The assertion of Lemma 2.5.2 holds.

Proof. Left as exercise. □

The universal property of henselization induces a map $A^b \rightarrow \widehat{A}$.

Proposition 2.A.9. *For a valued field (K, ν) , the following are equivalent:*

(a) K is henselian,

(b) every finite extension L of K admits a unique extension of the valuation ν .

Proof. Suppose that K is henselian. Given Lemma 2.A.8, we can repeat the proof of Proposition 2.5.3 word for word. The first paragraph of the proof of Theorem 2.5.1 shows that we can extend the valuation ring of K to L , which gives an extension of the valuation, easily seen to be unique. For the reverse direction, see [8, Theorem II 6.6]. □

Henselian rings will appear later in the course: the local ring $\mathcal{O}_{X,x}$ of a point x on a rigid analytic space X is not complete, but it is henselian.¹⁹

¹⁹ The same holds for complex analytic spaces, e.g. the local ring $\mathbb{C}\{t\}$ of power series with positive radius of convergence is henselian.

3

The Tate algebra

In this chapter, we fix a nonarchimedean field K . We denote by \mathcal{O} its valuation ring, by $k = \mathcal{O}/\mathfrak{m}$ its residue field, and by $t \in \mathfrak{m}$ a fixed pseudouniformizer.

We first introduce the Tate algebra, slightly emphasizing the “algebraic” point of view. We equip it with the Gauss norm, for which we give a geometric interpretation which facilitates the verification of some basic properties like multiplicativity or the Maximum Principle. The Gauss norm makes the Tate algebra into a Banach K -algebra; we prove that it satisfies a universal property in the category of Banach K -algebras. Next, we prove that the Tate algebra satisfies a number of favorable algebraic or topological properties, namely: ¹

- it satisfies a version of Noether normalization,
- it is Noetherian,
- all of its ideals are closed,
- the residue fields of its maximal ideals are finite extensions of K .

In the appendix, written jointly with Alex Youcis, we figure out one can view Banach spaces over K algebraically through the lens of \mathcal{O}/t^n -modules.

3.1 Definition of the Tate algebra

Definition 3.1.1. The *algebra of restricted power series* in r variables is the t -adic completion of the polynomial algebra $\mathcal{O}[X_1, \dots, X_r]$:

$$\mathcal{O}\langle X_1, \dots, X_r \rangle = \varprojlim_n \mathcal{O}[X_1, \dots, X_r]/(t^n) = \varprojlim_n ((\mathcal{O}/t^n)[X_1, \dots, X_r]).$$

The *Tate algebra* in r variables is the localization

$$K\langle X_1, \dots, X_r \rangle = \mathcal{O}\langle X_1, \dots, X_r \rangle \otimes_{\mathcal{O}} K = \mathcal{O}\langle X_1, \dots, X_r \rangle \left[\frac{1}{t} \right].$$

Let $\mathfrak{n} = (t, X_1, \dots, X_r) \subseteq \mathcal{O}[X_1, \dots, X_r]$. The \mathfrak{n} -adic completion of $\mathcal{O}[X_1, \dots, X_r]$ is the ring of formal power series

$$\mathcal{O}[[X_1, \dots, X_r]] = \varprojlim_n \mathcal{O}[X_1, \dots, X_r]/\mathfrak{n}^n.$$

¹ I mostly managed to avoid the rather tedious arguments using the Weierstrass Preparation Theorem and the theory of local and B -rings used in [4, Chapter 2]. Matter of taste, I guess.

Since $\mathfrak{n} \supseteq (t)$, we get the induced map on the respective completions:

$$\mathcal{O}\langle X_1, \dots, X_r \rangle \rightarrow \mathcal{O}[[X_1, \dots, X_r]]. \quad (3.1)$$

Lemma 3.1.2. *The map (3.1) is injective, and its image consists of the power series whose coefficients tend to zero:*²

$$\mathcal{O}\langle X_1, \dots, X_r \rangle \simeq \left\{ \sum_{\mathbf{n} \in \mathbf{N}^r} a_{\mathbf{n}} \mathbf{x}^{\mathbf{n}} \in \mathcal{O}[[X_1, \dots, X_r]] : a_{\mathbf{n}} \rightarrow 0 \text{ as } |\mathbf{n}| \rightarrow \infty \right\}.$$

Proof. We define the inverse homomorphism φ . Let $f = \sum a_{\mathbf{n}} \mathbf{X}^{\mathbf{n}} \in \mathcal{O}[[\mathbf{X}]]$ be an element of the right hand side. The condition that $a_{\mathbf{n}} \rightarrow 0$ means precisely that for every $m \geq 0$, all but finitely many of the coefficients $a_{\mathbf{n}}$ are divisible by t^m . Thus, for every $m \geq 0$, the image f_m of f in $\mathcal{O}[[\mathbf{X}]]/t^m = (\mathcal{O}/t^m)[[\mathbf{X}]]$ is a polynomial. The elements $f_m \in (\mathcal{O}/t^m)[[\mathbf{X}]]$ form a compatible system, and give rise to an element $\varphi(f)$ of $\mathcal{O}\langle \mathbf{X} \rangle$. One easily checks that φ is the inverse to (3.1). \square

By inverting t , we obtain an isomorphism

$$K\langle X_1, \dots, X_r \rangle \simeq \left\{ \sum_{\mathbf{n} \in \mathbf{N}^r} a_{\mathbf{n}} \mathbf{X}^{\mathbf{n}} \in K[[X_1, \dots, X_r]] : a_{\mathbf{n}} \rightarrow 0 \text{ as } |\mathbf{n}| \rightarrow \infty \right\}.$$

As we have observed in §1.1, the right hand side is precisely the algebra of power series with coefficients in K which converge in the unit disc

$$\mathbf{D}^r(K) = \{(x_1, \dots, x_r) \in K : |x_i| \leq 1 \text{ for } i = 1, \dots, r\}.$$

In particular, this implies that if for $f \in K[[X_1, \dots, X_r]]$ the series $f(\mathbf{x})$ converges for all $\mathbf{x} \in \mathbf{D}^r(K)$, then it also converges for all $\mathbf{x} \in \mathbf{D}^r(\overline{K})$.

3.2 The topology on $K\langle X_1, \dots, X_r \rangle$ and the Gauss norm

The ring $\mathcal{O}\langle X_1, \dots, X_r \rangle$, being defined as a completion, carries a natural inverse limit topology, called the *t-adic topology*. It extends uniquely to a topology of the Tate algebra $K\langle X_1, \dots, X_r \rangle$ for which $\mathcal{O}\langle X_1, \dots, X_r \rangle$ is an open subring; that topology can be described as the inductive limit topology, since

$$K\langle X_1, \dots, X_r \rangle = \bigcup_{n \geq 0} t^{-n} \mathcal{O}\langle X_1, \dots, X_r \rangle.$$

Below, we describe the natural norm inducing these topologies.

Definition 3.2.1. The *Gauss norm* on $K\langle X_1, \dots, X_r \rangle$ is defined by

$$|f| = \max\{|a_{\mathbf{n}}| : \mathbf{n} \in \mathbf{N}^r\} \quad \text{if } f = \sum_{\mathbf{n} \in \mathbf{N}^r} a_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}.$$

In other words, $|f|$ is the infimum of the values of $|c|$ for $c \in K^\times$ such that $c^{-1}f \in \mathcal{O}\langle X_1, \dots, X_r \rangle$. In particular, we have

$$\mathcal{O}\langle X_1, \dots, X_r \rangle = \{f \in K\langle X_1, \dots, X_r \rangle : |f| \leq 1\}.$$

The topology on $\mathcal{O}\langle X_1, \dots, X_r \rangle$ induced by the metric $d(x, y) = |x - y|$ is the *t-adic topology*.

² Here we use the multi-index notation: if $\mathbf{n} = (n_1, \dots, n_r) \in \mathbf{N}^r$, we set $\mathbf{X}^{\mathbf{n}} = X_1^{n_1} \cdots X_r^{n_r}$ and $|\mathbf{n}| = n_1 + \dots + n_r$.

Compare with Exercise 2 on Problem Set 2.

The geometric interpretation: suppose that K is discretely valued, and that $t \in \mathcal{O}$ is a uniformizer. Then $X = \text{Spec } \mathcal{O}[X_1, \dots, X_r] = \mathbf{A}_{\mathcal{O}}^r$ is a Noetherian regular scheme, and $Y = \{t = 0\} = \mathbf{A}_{\mathcal{O}}^r$ is a prime divisor on X . Therefore Y defines a valuation of height one v_Y on $k(X)$ (“order of zero or pole along Y ”). It agrees with the Gauss norm in the weak sense that for $f \in K[X_1, \dots, X_r] \subseteq K\langle X_1, \dots, X_r \rangle$, we have

$$|f|_{\text{Gauss}} = |t|^{-v_Y(f)}.$$

In fact, $K[X_1, \dots, X_r]$ is dense in $K\langle X_1, \dots, X_r \rangle$ with respect to the t -adic topology, and the Gauss norm is the unique continuous extension of the norm $|t|^{-v_Y(f)}$ to $K\langle X_1, \dots, X_r \rangle$.

The proofs of the following two easy results employ the above intuition.

Lemma 3.2.2 (The Gauss norm is multiplicative). *We have $|fg| = |f| \cdot |g|$ for $f, g \in K\langle X_1, \dots, X_r \rangle$.*

Proof. Clearly this holds if $f \in K$ is a constant. We can therefore rescale f and g so that $|f| = 1 = |g|$. Equivalently $f, g \in \mathcal{O}\langle X_1, \dots, X_r \rangle$ and their residues modulo the maximal ideal $\mathfrak{m} \subseteq \mathcal{O}$

$$\bar{f}, \bar{g} \in \mathcal{O}\langle X_1, \dots, X_r \rangle / \mathfrak{m} = k[X_1, \dots, X_r]$$

are nonzero. Since $k[X_1, \dots, X_r]$ is a domain, $fg \in \mathcal{O}\langle X_1, \dots, X_r \rangle$ has nonzero image $\bar{f}\bar{g}$ in $k[X_1, \dots, X_r]$, and hence $|fg| = 1 = |f| \cdot |g|$. \square

Proposition 3.2.3 (The Maximum Principle). *For $f \in K\langle X_1, \dots, X_r \rangle$, we have*

$$|f| = \sup \left\{ |f(x_1, \dots, x_r)| : (x_1, \dots, x_r) \in \bar{K}^r, |x_i| \leq 1 \right\}.$$

Proof. As in the previous proof, we can reduce to the case $|f| = 1$.

Clearly, the right hand side is ≤ 1 ; we will show it equals 1. We have $f \in \mathcal{O}\langle X_1, \dots, X_r \rangle$ and its image $\bar{f} \in k[X_1, \dots, X_r]$ is nonzero. We can therefore find a point $(\bar{\xi}_1, \dots, \bar{\xi}_r) \in \bar{k}^r$ such that $\bar{f}(\bar{\xi}_1, \dots, \bar{\xi}_r) \neq 0$. Now \bar{k} is the residue field of (the integral closure of \mathcal{O} in) \bar{K} ; let $(\xi_1, \dots, \xi_r) \in \bar{K}^r$ be an element lifting $(\bar{\xi}_1, \dots, \bar{\xi}_r)$. Then $|\xi_i| \leq 1$ and $|f(\xi_1, \dots, \xi_r)| = 1$. \square

Remark 3.2.4. The above proof shows three things in addition. First, the supremum is a maximum, and therefore attained in L^r for L a finite extension of K . Second, if the residue field \bar{k} is infinite, the above maximum is attained at a point in K^r . Lastly, the maximum is attained at a point with $|x_1| = \dots = |x_r| = 1$.

The Gauss norm makes the Tate algebra into a Banach K -algebra, as defined below.

Definition 3.2.5 (Banach spaces and Banach algebras). Let V be a vector space over K . A *vector space norm* on V is a function

$$|\cdot|: V \rightarrow [0, \infty)$$

such that

- i. $|xv| = |x| \cdot |v|$ for $x \in K, v \in V$,

- ii. $|v| = 0$ if and only if $v = 0$,
- iii. $|v + w| \leq \max\{|v|, |w|\}$ for $v, w \in V$.

It is called a *Banach norm* if V is complete with respect to the induced metric $d(x, y) = |x - y|$. A *Banach space* over K is a vector space over K equipped with a Banach norm.

Let A be a K -algebra. A *K -algebra norm* on A is a vector space norm $|\cdot|$ on A which satisfies

- iv. $|vw| \leq |v| \cdot |w|$ for $v, w \in A$.

It is a *Banach algebra norm* if $|\cdot|$ is a Banach norm. A *Banach K -algebra* is a K -algebra over K equipped with a Banach norm.

A linear map $f: V \rightarrow W$ between Banach spaces over K is continuous if and only if it is *bounded* in the sense that $|f(v)| \leq C \cdot |v|$ ($v \in V$) for some constant C independent of v . This implies in particular that a continuous $f: V \rightarrow W$ is uniformly continuous.

Proposition 3.2.6. *The Tate algebra $K\langle X_1, \dots, X_r \rangle$ is a Banach algebra when equipped with the Gauss norm.*

Proof. Axioms i.–iii. are clear, and iv. follows from Lemma 3.2.2. It remains to show that $K\langle X_1, \dots, X_r \rangle$ is complete. It suffices to show that the closed unit ball $\{|f| \leq 1\} = \mathcal{O}\langle X_1, \dots, X_r \rangle$ is complete. This in turn follows from the fact that $\mathcal{O}\langle X_1, \dots, X_r \rangle$ is t -adically complete. \square

Corollary 3.2.7. *The Tate algebra $K\langle X_1, \dots, X_r \rangle$ is the completion of $K[X_1, \dots, X_r]$ with respect to the Gauss norm.*

Proof. It suffices to observe that $\mathcal{O}[X_1, \dots, X_r]$ is dense in $\mathcal{O}\langle X_1, \dots, X_r \rangle$, which follows from the definition (and the fact that the metric topology induced by the Gauss norm agrees with the t -adic topology). \square

3.3 The universal property

Definition 3.3.1. Let A be a Banach K -algebra. An element $a \in A$ is *powerbounded* if the set $\{a^n : n \geq 1\}$ is bounded, meaning that $\{|a^n| : n \geq 1\}$ is bounded from above. We denote the set of powerbounded elements by $A^\circ \subseteq A$.

The subset $A^\circ \subseteq A$ is a subring. If the norm on A is multiplicative, then $a \in A^\circ$ if and only if $|a| \leq 1$; therefore $A^\circ = \{|a| \leq 1\}$ is an open subring. Thus for $A = K\langle X_1, \dots, X_r \rangle$ we have $A^\circ = \mathcal{O}\langle X_1, \dots, X_r \rangle$.

Every continuous homomorphism $A \rightarrow B$ maps A° into B° . Since the element $X \in K\langle X \rangle$ is powerbounded, for every Banach K -algebra we obtain a map

$$\varphi \mapsto \varphi(X) \quad : \quad \text{Hom}_K(K\langle X \rangle, A) \rightarrow A^\circ, \quad (3.2)$$

where for Banach K -algebras A and B , $\text{Hom}_K(B, A)$ denotes the set of all *continuous* K -algebra homomorphisms $B \rightarrow A$.

Warning. If A is not reduced, then the subring A° is not very well-behaved.

For example, if $A = K\langle X \rangle/(X^2)$ then $A^\circ = \mathcal{O} \oplus K \cdot X$ is neither bounded nor t -adically separated.

Lemma 3.3.2. *The maps (3.2) are bijective and define an isomorphism between the functors $A \mapsto \text{Hom}_K(K\langle X \rangle, A)$ and $A \mapsto A^\circ$ from Banach K -algebras to sets. In other words, $K\langle X \rangle$ represents the functor $A \mapsto A^\circ$.*

Proof. Since $K[X]$ is dense in $K\langle X \rangle$ (Corollary 3.2.7), any two continuous K -algebra homomorphisms $\varphi, \psi: K\langle X \rangle \rightarrow A$ with $\varphi(X) = \psi(X)$ have to coincide. This shows injectivity. To show that $\varphi \mapsto \varphi(X)$ is surjective, let $a \in A^\circ$ and let $\varphi: K[X] \rightarrow A$ be the unique K -algebra homomorphism sending X to a . To extend φ to the completion $K\langle X \rangle$ of $K[X]$ with respect to the Gauss norm, it suffices to show that φ is (uniformly) continuous, i.e. that

$$|\varphi(f)| \leq C \cdot |f| \quad \text{for some } C > 0.$$

Since a is powerbounded, there exists a C such that $|a^n| \leq C$ for all $n \geq 0$. But then, for $f = \sum_{i=0}^m b_i X^i \in K[X]$, we have

$$|\varphi(f)| = \left| \sum_{i=0}^m b_i a^i \right| \leq \max\{|b_i|\} \cdot \max\{|a^n|\} \leq |f| \cdot C. \quad \square$$

3.4 The Tate algebra is Noetherian

The goal of this section is to prove that $K\langle X_1, \dots, X_r \rangle$ is Noetherian.

Proposition 3.4.1 (Warm-up). *Suppose that K is discretely valued, i.e. \mathcal{O} is a dvr. Then $\mathcal{O}\langle X_1, \dots, X_r \rangle$ and $K\langle X_1, \dots, X_r \rangle$ are Noetherian.*

Proof. Since \mathcal{O} is Noetherian, so is the polynomial algebra $\mathcal{O}[X_1, \dots, X_r]$. The completion of a Noetherian ring with respect to an ideal is Noetherian [2, Theorem 10.26], thus $\mathcal{O}\langle X_1, \dots, X_r \rangle$ is Noetherian. Finally, the localization of a Noetherian ring is Noetherian, and therefore $K\langle X_1, \dots, X_r \rangle$ is Noetherian as well. \square

However, if the valuation is nondiscrete, then \mathcal{O} will not be Noetherian: the maximal ideal is not finitely generated, in fact it satisfies $\mathfrak{m} = \mathfrak{m}^2$. Thus $\mathcal{O}\langle X_1, \dots, X_r \rangle$ is non-Noetherian as well, for the same reason. That reason disappears when we invert t .

The proof below loosely follows Tian's notes [11], with some simplifications.

Proposition 3.4.2 (Noether normalization). *Let $I \subseteq K\langle X_1, \dots, X_r \rangle$ be a closed ideal.³ Then there exists a finite and injective K -algebra homomorphism*

$$K\langle Y_1, \dots, Y_s \rangle \hookrightarrow K\langle X_1, \dots, X_r \rangle / I \quad \text{for some } s \leq r.$$

Proof. The idea of the proof is to deduce the statement from the usual Noether normalization lemma over k . We shall use the algebra $\mathcal{O}\langle X_1, \dots, X_r \rangle$ as an intermediary between the Tate algebra $K\langle X_1, \dots, X_r \rangle$ and the polynomial ring $k[X_1, \dots, X_r]$.

Let $J = I \cap \mathcal{O}\langle X_1, \dots, X_r \rangle$ and $B = \mathcal{O}\langle X_1, \dots, X_r \rangle / J$. Note that J is open in I , we have $I = J \cdot K\langle X_1, \dots, X_r \rangle$, and J is closed in $\mathcal{O}\langle X_1, \dots, X_r \rangle$. The last fact implies that

$$B \simeq \varprojlim_n B/t^n, \quad B/t^n = (\mathcal{O}/t^n)[X_1, \dots, X_r]/J.$$

Similarly, $K\langle X_1, \dots, X_r \rangle$ represents the functor $A \mapsto (A^\circ)^r$.

³ We shall soon prove that every ideal in $K\langle X_1, \dots, X_r \rangle$ is closed.

Noether normalization applied to $B/\mathfrak{m} = k[X_1, \dots, X_r]/J$ produces a finite injective map

$$k[Y_1, \dots, Y_s] \rightarrow B/\mathfrak{m}$$

which we can lift to a map $\mathcal{O}\langle Y_1, \dots, Y_s \rangle \rightarrow B$. Indeed, we can certainly lift it to an \mathcal{O} -algebra map $\mathcal{O}[Y_1, \dots, Y_r] \rightarrow B$, and upon taking t -adic completion we obtain the desired $\mathcal{O}\langle Y_1, \dots, Y_s \rangle \rightarrow B$ (because B is t -adically complete). We want to show that the latter map is finite and injective as well.

Injectivity is easy: let $f \in \mathcal{O}\langle Y_1, \dots, Y_s \rangle$ and write $f = cg$ with $c \in \mathcal{O}$ and $|g| = 1$. Then g has nonzero image in $k[Y_1, \dots, Y_s]$, and hence its image in B/\mathfrak{m} is nonzero. Since B is \mathcal{O} -torsion free (being a submodule of the K -module $K\langle X_1, \dots, X_r \rangle/I$), we see that f maps to zero only for $c = 0$.

For finiteness, as an intermediate step we will show that

$$\mathcal{O}/t[Y_1, \dots, Y_s] \rightarrow B/t$$

is finite. It suffices to show that the images of X_i in B/t are integral over $\mathcal{O}/t[Y_1, \dots, Y_s]$. Since their images in B/\mathfrak{m} are integral over $k[Y_1, \dots, Y_s]$, there exist monic polynomials $P_i \in \mathcal{O}\langle Y_1, \dots, Y_s \rangle[X]$ with $P_i(X_i) \in \mathfrak{m}B$. But then for $N \gg 0$ we have $P_i^N(X_i) \in tB$, i.e. the X_i are integral over $\mathcal{O}/t[Y_1, \dots, Y_s]$.

Now, let $\{Z_\alpha\}$ be a finite set of elements of B which generate B/t as a $\mathcal{O}/t[Y_1, \dots, Y_s]$ -module. Fix $W_0 \in B$ and write

$$\begin{aligned} W_0 &= \sum_{\alpha} f_{0,\alpha} Z_{\alpha} + t W_1 \\ &= \sum_{\alpha} (f_{0,\alpha} + t f_{1,\alpha}) Z_{\alpha} + t^2 W_2 \\ &= \dots \stackrel{?}{=} \sum_{\alpha} f_{\alpha} Z_{\alpha} \end{aligned}$$

where $f_{\alpha} = \sum_n f_{n,\alpha} t^n$. Indeed, the difference of the two sides of $\stackrel{?}{=}$ belongs to $\bigcap_n t^n B = 0$. Therefore Z_{α} generate B over $\mathcal{O}\langle Y_1, \dots, Y_s \rangle$.⁴ \square

Remark 3.4.3. The above proof shows that we can choose the finite injective map so that it factors through $K\langle X_1, \dots, X_r \rangle$. This is not automatic, for example in the situation below

$$\begin{array}{ccc} & & K\langle X \rangle \\ & \nearrow & \downarrow \\ K\langle Y \rangle & \xrightarrow{Y \mapsto t^{-1}X} & K\langle X \rangle/(X^2) \end{array}$$

there does not exist a dotted arrow making the triangle commute. Indeed, the element $t^{-1}X \in A$ is nilpotent and hence power-bounded, but it cannot be lifted to a power-bounded element of $K\langle X \rangle$. We will need this observation in §4.2, where the above issue will be clarified.

Proposition 3.4.4. *The Tate algebra $K\langle X_1, \dots, X_r \rangle$ is Noetherian.*

Proof. We prove this by induction on r . Let $I \subseteq K\langle X_1, \dots, X_r \rangle$ be a nonzero ideal. Pick $f \in I$ with $|f| = 1$. It is enough to show that

⁴ The argument presented in the final paragraph shows more generally that if A is a t -adically complete \mathcal{O} -algebra and M is a t -adically separated A -module, then elements $e_1, \dots, e_n \in M$ which generate M/t also generate M (“ t -adic Nakayama’s lemma”).

$K\langle X_1, \dots, X_r \rangle / (f)$ is Noetherian, for then the image $I/(f)$ is finitely generated and hence so is I .

The ideal (f) is closed, as multiplication by f

$$f : K\langle X_1, \dots, X_r \rangle \rightarrow K\langle X_1, \dots, X_r \rangle$$

is an isometry onto its image (f) . We can therefore apply Noether normalization (Proposition 3.4.2) to obtain a finite and injective homomorphism

$$K\langle Y_1, \dots, Y_s \rangle \hookrightarrow K\langle X_1, \dots, X_r \rangle / (f).$$

Moreover, since $|f| = 1$, we must have $s < r$ by construction. By induction, $K\langle Y_1, \dots, Y_s \rangle$ is Noetherian and hence so is $K\langle X_1, \dots, X_r \rangle / (f)$. \square

Proposition 3.4.5. *Every ideal in $K\langle X_1, \dots, X_r \rangle$ is closed.*

Proof. Let $I \subseteq K\langle X_1, \dots, X_r \rangle$ be an ideal and let \bar{I} be its closure. Then \bar{I} , again an ideal, is finitely generated: $\bar{I} = (f_1, \dots, f_s)$. Using the density of I in \bar{I} , we will show that we can find another system of generators (g_1, \dots, g_s) with $g_i \in I$, showing $I = \bar{I}$.

Consider the surjective and bounded map of Banach spaces

$$K\langle X_1, \dots, X_r \rangle^{\oplus s} \rightarrow \bar{I}, \quad (h_1, \dots, h_s) \mapsto \sum b_i f_i.$$

By the Open Mapping Theorem⁵, there exists a $C > 0$ such that for every $f \in \bar{I}$ there exist $h_1, \dots, h_s \in K\langle X_1, \dots, X_r \rangle$ with $f = \sum b_i f_i$ and $|h_i| \leq C \cdot |f|$.

Since $I \subseteq \bar{I}$ is dense, we can find $g_1, \dots, g_s \in I$ with $|g_i - f_i| < C^{-1}$. By the previous paragraph, there exist $h_{ij} \in K\langle X_1, \dots, X_r \rangle$ ($1 \leq i, j \leq s$) such that

$$g_i - f_i = \sum_j h_{ij} f_j \quad \text{and} \quad |h_{ij}| < 1.$$

Rewrite this as

$$g_i = \sum_j H_{ij} f_j, \quad H_{ij} = h_{ij} + \delta_{ij},$$

so that the matrix $H = [H_{ij}]$ satisfies $|H - \text{Id}| < 1$ (for the supremum norm on matrix entries). It is easy to see (see Problem 2 on PS3) that this implies that H is invertible, showing $\bar{I} = (f_1, \dots, f_s) \subseteq (g_1, \dots, g_s) \subseteq I$. \square

3.5 Maximal ideals

Recall that by Nullstellensatz, for an algebraically closed field k , the maximal ideals in $k[X_1, \dots, X_r]$ are in bijection with k^r . If k is not necessarily algebraically closed, and \bar{k} is an algebraic closure, then maximal ideals in $k[X_1, \dots, X_r]$ correspond to orbits of the action of the Galois group $\text{Gal}(\bar{k}/k)$ on \bar{k}^r . The case of the Tate algebra is similar.

Proposition 3.5.1. *There is a bijection between the set $\text{Max} K\langle X_1, \dots, X_r \rangle$ of maximal ideals in $K\langle X_1, \dots, X_r \rangle$ and the set of orbits of the action of the Galois group $\text{Gal}(\bar{K}/K)$ on*

$$\mathbf{D}^r(\bar{K}) = \{(x_1, \dots, x_r) \in \bar{K}^r : |x_i| \leq 1\},$$

where $|\cdot|$ is the unique extension of the norm on K to \bar{K} .

⁵ **Open Mapping Theorem.** *A surjective continuous map $\pi : V \rightarrow W$ of Banach spaces over K is open. That is, there exists a $C > 0$ such that $\{|w| \leq 1\}$ is contained in $\pi(\{|v| \leq C\})$.*

Proof. Open your Functional Analysis textbook and check that the proof works without change in the non-Archimedean setting. \square

Proof. For $x = (x_1, \dots, x_r) \in \mathbf{D}^r(\overline{K})$, let

$$\mathfrak{m}_x = \{f \in K\langle X_1, \dots, X_r \rangle : f(x) = 0\}$$

(note that $f(x)$ makes sense because $|x_i| \leq 1$). This is a maximal ideal, as the image of the evaluation map

$$f \mapsto f(x): K\langle X_1, \dots, X_r \rangle \rightarrow \overline{K}$$

is a subring of \overline{K} containing K and hence is a field. Moreover, Galois conjugate points give the same ideal, so we get a map $x \mapsto \mathfrak{m}_x$ from one side to the other.

Conversely, let \mathfrak{n} (the notation \mathfrak{m} already being reserved for the maximal ideal in \mathcal{O}) be a maximal ideal in $K\langle X_1, \dots, X_r \rangle$. Applying Noether normalization, we see that the residue field $L = K\langle X_1, \dots, X_r \rangle/\mathfrak{n}$ is finite over $K\langle X_1, \dots, X_s \rangle$ for some s . But this implies that the latter ring is a field, so $s = 0$ and L is a finite extension of K . Embedding it into \overline{K} , we obtain a homomorphism

$$\varphi: K\langle X_1, \dots, X_s \rangle \rightarrow L \rightarrow \overline{K}.$$

Let $x_i = \varphi(X_i) \in \overline{K}$. Thus x_i are powerbounded, and hence $|x_i| \leq 1$. This gives a point $x = (x_1, \dots, x_r) \in \mathbf{D}^r(\overline{K})$, well-defined up to the choice of the embedding of L in \overline{K} . This gives a map $\mathfrak{n} \mapsto x$ in the other direction.

As such embeddings are permuted by the Galois group, it is clear that $\mathfrak{m}_x \mapsto x$. If $\mathfrak{n} \mapsto x$, then $\mathfrak{n} \subseteq \mathfrak{m}_x$, and hence they are equal since both are maximal. We have thus constructed mutually inverse bijections. \square

Corollary 3.5.2. *Every K -algebra homomorphism*

$$K\langle Y_1, \dots, Y_s \rangle \rightarrow K\langle X_1, \dots, X_r \rangle$$

is continuous.

Proof. By the Maximum Principle (Proposition 3.2.3), the Gauss norm on $K\langle X_1, \dots, X_r \rangle$ agrees with the *supremum norm*

$$|f|_{\text{sup}} = \sup\{|f \bmod \mathfrak{n}| : \mathfrak{n} \in \text{Max} K\langle X_1, \dots, X_r \rangle\},$$

where $|f \bmod \mathfrak{n}|$ is the norm of the image of f in the residue field $L = K\langle X_1, \dots, X_r \rangle/\mathfrak{n}$. This definition of the Gauss norm is *intrinsic* to the K -algebra structure on $K\langle X_1, \dots, X_r \rangle$. It is also straightforward to check using $|\cdot|_{\text{Gauss}} = |\cdot|_{\text{sup}}$ that for every K -algebra homomorphism

$$\varphi: K\langle Y_1, \dots, Y_s \rangle \rightarrow K\langle X_1, \dots, X_r \rangle$$

we have $|\varphi(f)| \leq |f|$, i.e. f is not only continuous but even *contracting*. \square

3.6 More commutative algebra

We state the following additional results without giving a proof.

Theorem 3.6.1. (a) *The Tate algebra is Jacobson (every prime ideal is the intersection of maximal ideals).*

See [4, Proposition 2.2/16].

(b) *The Tate algebra is regular, of Krull dimension n , and excellent.*

See [5, §1.1] and references therein.

(c) *Every ideal $I \subseteq K\langle X_1, \dots, X_r \rangle$ admits a system of generators (f_1, \dots, f_s) with $|f_i| = 1$ and such that every $f \in I$ we can write $f = \sum f_i g_i$ with $|g_i| \leq |f|$*

See [4, Corollary 2.3/7].

3.A Banach spaces (with Alex Youcis)

Still slightly incomplete.

The goal of this slightly persnickety appendix, only tangentially related to the lecture, is to explicate the notion of a Banach space over K in terms of \mathcal{O}/t^n -modules. The main result (Proposition 3.A.10) describes the category Ban_K of Banach spaces over K as a localization of the category $\text{Mod}_{\mathcal{O}}^{\wedge}$ of *complete* \mathcal{O} -modules (which itself is the inverse limit of the categories $\text{Mod}_{\mathcal{O}/t^n}$) with respect to *topological isogenies*, i.e. morphisms whose kernel and cokernel have dense torsion submodules.

As before, we work over a non-Archimedean field K , denote by $\mathcal{O} \subseteq K$ be its valuation ring, and fix a pseudouniformizer $t \in \mathcal{O}$.

3.A.1 Torsion-free \mathcal{O} -modules

Mod_A for a ring A is the category of all A -modules, and Mod_A^f is the full subcategory of flat A -modules.

For $M \in \text{Mod}_{\mathcal{O}}$, we define its *torsion submodule*

$$M_{\text{tors}} = \bigcup_{n \geq 0} \ker(t^n: M \rightarrow M).$$

The module M is *torsion* (resp. *torsion-free*) if $M_{\text{tors}} = M$ (resp. $M_{\text{tors}} = 0$). We have the following basic result:

Lemma 3.A.1. *An \mathcal{O} -module M is flat if and only if it is torsion-free.*

Since the module M/M_{tors} is torsion-free, we have a functorial way of making any given \mathcal{O} -module flat. Since every map $M \rightarrow N$ where N is torsion-free has to map M_{tors} to zero, we obtain:

Lemma 3.A.2. *The functor*

$$M \mapsto M/M_{\text{tors}}: \text{Mod}_{\mathcal{O}} \rightarrow \text{Mod}_{\mathcal{O}}^f$$

is a left adjoint to the inclusion $\text{Mod}_{\mathcal{O}}^f \subseteq \text{Mod}_{\mathcal{O}}$.

3.A.2 Complete \mathcal{O} -modules

The *completion* of an \mathcal{O} -module M is the inverse limit

$$\widehat{M} = \varprojlim_n M/t^n M.$$

A \mathcal{O} -module M is *complete* if the natural map $M \rightarrow \widehat{M}$ is an isomorphism. We denote by $\text{Mod}_{\mathcal{O}}^{\wedge}$ the full subcategory of $\text{Mod}_{\mathcal{O}}$ consisting of complete \mathcal{O} -modules. The completion functor

$$M \mapsto \widehat{M}: \text{Mod}_{\mathcal{O}} \rightarrow \text{Mod}_{\mathcal{O}}^{\wedge}$$

is a left adjoint to the inclusion $\text{Mod}_{\mathcal{O}}^{\wedge} \subseteq \text{Mod}_{\mathcal{O}}$.

We denote by $\text{Mod}_{\mathcal{O}}^{\wedge, f}$ the full subcategory of flat and complete \mathcal{O} -modules. The completion of a flat \mathcal{O} -module is again flat, and again the completion functor $\text{Mod}_{\mathcal{O}}^f \rightarrow \text{Mod}_{\mathcal{O}}^{\wedge, f}$ is a left adjoint to the inclusion functor.

We have equivalences of categories

$$\mathrm{Mod}_\theta^\wedge = 2\text{-}\varprojlim_n \mathrm{Mod}_{\theta/t^n} \quad \text{and} \quad \mathrm{Mod}_\theta^{\wedge, f} = 2\text{-}\varprojlim_n \mathrm{Mod}_{\theta/t^n}^f,$$

where for an inverse system of categories $(\mathcal{C}_n, \pi_n: \mathcal{C}_{n+1} \rightarrow \mathcal{C}_n)$, we define its 2-categorical inverse limit $2\text{-}\varprojlim_n \mathcal{C}_n$ as consisting of systems of objects and isomorphisms $(x_n \in \mathcal{C}_n, \iota_n: \pi_n(x_{n+1}) \simeq x_n)$, and where morphisms are systems of maps $(x'_n \rightarrow x_n)$ commuting with the maps ι'_n, ι_n .

Warning: The category $\mathrm{Mod}_\theta^\wedge$ has kernels and cokernels. The kernel is simply the kernel in Mod_θ , and the cokernel is the completion of the usual cokernel. However, $\mathrm{Mod}_\theta^\wedge$ is not abelian. The reason for that is that the image of a map need not be closed.

See [10, Tag 07JQ].

Lemma 3.A.3. *The functor $M \mapsto (M/M_{\mathrm{tors}})^\wedge$ is a left adjoint to the inclusion $\mathrm{Mod}_\theta^{\wedge, f} \subseteq \mathrm{Mod}_\theta^\wedge$.*

Before we begin, we start with the following ancillary lemma:

Lemma 3.A.4. *Let M be an object of $\mathrm{Mod}_\theta^\wedge$ and N a subspace of M . Then, there is a natural embedding*

$$N/\overline{M} \rightarrow (N/M)^\wedge$$

with dense image.

Proof. Let us note that

$$\begin{aligned} (N/M)^\wedge &= \varprojlim_n (M/N)/t^n(M/N) \\ &= \varprojlim_n M/(t^n, N) \end{aligned}$$

So, let us then observe that we have a natural map

$$M \rightarrow \varprojlim_n M/(t^n, N)$$

We claim that the kernel of this map is precisely \overline{N} . Indeed, to show that \overline{N} is in the kernel we need to show that \overline{N} projects to zero in (t^n, N) for every n . But, take x in \overline{N} and write $x = \lim y_n$ with y_n in N for all n and $x - y_n \in t^n M$. Then, evidently x projects to 0 in $M/(t^n, N)$ since x is in $y_n + t^n M \subseteq (t^n, M)$. Conversely, suppose that x maps to zero in $\varprojlim_n M/(t^n, N)$. Then, by definition, for all $n \geq 0$ we have that we can write $x = y_n + t^n z_n$ for some y_n in N and z_n in M . In particular, from this we see that $x = \lim y_n$ and thus x is in \overline{N} .

From this we see that we get an injection

$$M/\overline{N} \rightarrow \varprojlim_n M/(t^n, N) = (M/N)^\wedge$$

To see that it has dense image it suffices to note that for all n we have the composition

$$M/\overline{N} \rightarrow \varprojlim_n M/(t^n, N) \rightarrow M/(t^n, N)$$

is surjective, from where the claim follows. \square

From this we deduce the following:

Corollary 3.A.5. *Let M be an object of $\mathrm{Mod}_\theta^\wedge$. Then, M_{tors} is dense in M if and only if $(M/M_{\mathrm{tors}})^\vee$ is zero.*

We now return to Lemma 3.A.3:

Proof of Lemma 3.A.3. We need to show that for every object M of $\text{Mod}_{\mathcal{O}}^{\wedge}$ and every object N of $\text{Mod}_{\mathcal{O}}^{\wedge, f}$ we have that the natural bijection

$$\text{Hom}((M/M_{\text{tors}})^{\vee}, N) \cong \text{Hom}(M, N)$$

But, we note that evidently the natural map

$$\text{Hom}(M, N) \rightarrow \text{Hom}(M/M_{\text{tors}}, N)$$

is a bijection since N is \mathcal{O} -flat. Moreover, since N is \mathcal{O} -complete we have that the natural map

$$\text{Hom}(M/M_{\text{tors}}, N) \rightarrow \text{Hom}((M/M_{\text{tors}})^{\wedge}, N)$$

is a bijection. The claim follows. \square

3.A.3 Banach spaces

See Definition 3.2.5 for the definition of a Banach space. A linear map $f: V \rightarrow W$ between Banach spaces over K is called *bounded* if there exists a $c \in [0, \infty)$ such that

$$|f(v)| \leq c|v| \quad \text{for all } v \in V.$$

We denote by $\text{Hom}(V, W)$ the linear space of such maps. It is stable under composition, and we denote the category of all Banach K -spaces and bounded maps by Ban_K .

We then have the following well-known result (e.g. see [3, §2.1.6] and [3, §2.1.8]):

Lemma 3.A.6. *Let V and W be Banach K -spaces. Then, a K -linear map $f: V \rightarrow W$ is bounded if and only if it's continuous. Moreover, the function*

$$|f| := \sup_{\substack{x \neq 0 \\ |x|=1}} \frac{|f(x)|}{|x|}$$

is a norm on $\text{Hom}(V, W)$ which endows $\text{Hom}(V, W)$ with the structure of a Banach K -space. Moreover, the following properties hold:

1. $|f| = \sup_{\substack{x \in V \\ |x|=1}} |f(x)|$
2. $|f(x)| \leq |f||x|$ for all x in V .
3. $|f \circ g| \leq |f||g|$ for any continuous map of Banach K -spaces $g: W \rightarrow U$.

3.A.4 Lattices

For $V \in \text{Ban}_K$, we write $V_0 = \{|v| \leq 1\}$. We then have the following elementary observation:

Lemma 3.A.7. *The subset V_0 is an \mathcal{O} -submodule which is \mathcal{O} -flat, complete, and such that the induced map $V_0 \otimes_{\mathcal{O}} K \rightarrow V$ is an isomorphism.*

Proof. Since $|xv| \leq |x||v|$ for all x in K and v in V we evidently see that V_0 is an \mathcal{O} -submodule of V . Since V is a K -module we know that it's \mathcal{O} -torsionfree and thus a fortiori the same holds true for V_0 which implies that it's \mathcal{O} -flat. Finally, we note that the induced map $V_0 \otimes_{\mathcal{O}} K \rightarrow V$ is an isomorphism as follows. Since K is \mathcal{O} -flat we have that the induced map $V_0 \otimes_{\mathcal{O}} K \rightarrow V \otimes_{\mathcal{O}} K$ is injective. But, we note that $V \otimes_{\mathcal{O}} K \cong V$ via the map which maps $v \otimes x$ to xv . Thus, we see that the induced map $V_0 \otimes_{\mathcal{O}} K \rightarrow V$ is an isomorphism if and only if for all v in V one can write $v = xv_0$ with x in K and v_0 in V_0 . But, this is clear since if $t^n v$ converges to 0 and so, since V_0 is open in V , must be in V_0 for some $n \geq 0$. We then can write $v = t^{-n}(t^n v)$. \square

If $f: V \rightarrow W$ is a continuous map of Banach K -spaces, then for $c \in K$ we have $f(V_0) \subseteq cW_0$ if and only if $|c| \geq |f|$. In particular, we see that if we set

$$\text{Hom}_0(V, W) := \{f \in \text{Hom}(V, W) : |f| \leq 1\}$$

then we have the equality

$$\text{Hom}_0(V, W) = \{f \in \text{Hom}(V, W) : f(V_0) \subseteq W_0\}$$

We define the category $\text{Ban}_{\mathcal{O}}$ to be the subcategory of Ban_K with the same underlying class of objects but where for V and W Banach K -spaces we set

$$\text{Hom}_{\text{Ban}_{\mathcal{O}}}(V, W) := \text{Hom}_0(V, W)$$

and call it the category of *Banach lattices*.

Proposition 3.A.8. *The functors*

$$-\otimes K : \text{Mod}_{\mathcal{O}}^{\wedge, f} \rightarrow \text{Ban}_{\mathcal{O}}, \quad V \mapsto V_0 : \text{Ban}_{\mathcal{O}} \rightarrow \text{Mod}_{\mathcal{O}}^{\wedge, f}$$

are mutually inverse equivalences of categories.

Before we explain the proof of this lemma, we remark as to how for M an object of $\text{Mod}_{\mathcal{O}}^{\wedge, f}$ we are regarding $M \otimes_{\mathcal{O}} K$ as a Banach K -space. Namely, we have the following simple observation:

Lemma 3.A.9. *Let M be an object of $\text{Mod}_{\mathcal{O}}^{\wedge, f}$. Then, the function*

$$|v| := \inf_{\substack{x \in K^{\times} \\ x^{-1}v \in M}} |x|$$

defines the structure of a Banach K -space on $M \otimes_{\mathcal{O}} K$. Moreover, if $f: M \rightarrow N$ is an \mathcal{O} -module map, then the induced map $f: M \otimes_{\mathcal{O}} K \rightarrow N \otimes_{\mathcal{O}} K$ is continuous.

Let us note that we are using the \mathcal{O} -flatness of M to regard M as a subgroup of $M \otimes_{\mathcal{O}} K$.

Proof of Lemma 3.A.9. Let us first verify that $|\cdot|$ really is a norm on $M \otimes_{\mathcal{O}} K$.

We first observe that if $|v| = 0$ then we have that $x^{-1}v$ is in M for all x in K^{\times} . From this it's easy to see that v is an element of M . Moreover, we see that we in fact have that v is an element of $t^n M$ for all $n \geq 0$. Since M is complete this implies that v is zero as desired.

To see that $|v + w| \leq \max(|v|, |w|)$ for all v and w in $M \otimes_{\theta} K$ is easy. Indeed, we note that if $x^{-1}v$ and $x^{-1}w$ are both in M then so then is $x^{-1}(v + w)$ from where the claim follows.

Finally, we show that for all x in K and v in $M \otimes_{\theta} K$ we have that $|xv| = |x||v|$. To see this, we merely note that

$$\begin{aligned} |xv| &= \inf_{\substack{y \in K^{\times} \\ y^{-1}xv \in M}} |y| \\ &= \inf_{\substack{y \in K^{\times} \\ y^{-1}v \in M}} |yx| \\ &= |x| \inf_{\substack{y \in K^{\times} \\ y^{-1}v \in M}} |y| \\ &= |x||v| \end{aligned}$$

as desired.

To see that $M \otimes_{\theta} K$ is complete is clear. Indeed, suppose that $\{v_n\}$ is a Cauchy sequence in $M \otimes_{\theta} K$. Let us note then that there exists some $n_0 \geq 0$ such that $|v_n - v_m| \leq 1$ for $n \geq n_0$. One then sees from the ultrametric inequality that $v_n - v_{n_0}$ is in M for all $n \geq n_0$. Then, we see that $v_n - v_{n_0}$ is a Cauchy sequence in M and thus, by the completeness of M , converges.

To see the claim concerning maps we proceed as follows. We need to show that $\lim f(x_n) = f(\lim x_n)$. Note though that if $\lim x_n = x$ then this implies $\lim(x_n - x) = 0$. Thus, we see that for all $N \geq 0$ there exists an n_0 such that for $n \geq n_0$ we have that $x_n - x \in t^N M$. We see then that $f(x_n - x) \in f(t^N M) \subseteq t^N N$. Thus, we see that $\lim f(x_n - x) = 0$ and thus $\lim f(x_n) = f(x)$ as desired. \square

We are now ready to prove Proposition 3.A.8:

Proof. (of Proposition 3.A.8) It suffices to show that $- \otimes_{\theta} K$ and $(-)_0$ are fully faithful and that they are inverses on isomorphism classes.

But, we note that for M and N objects of $\text{Mod}_{\theta}^{\wedge, f}$ that

$$\begin{aligned} \text{Hom}_0(M \otimes_{\theta} K, N \otimes_{\theta} K) &= \{f \in \text{Hom}(M \otimes_{\theta} K, N \otimes_{\theta} K) : f(M) \subseteq N\} \\ &= \text{Hom}(M, N) \end{aligned}$$

where this last map is an equality since as we observed in Lemma 3.A.9 the localization of any map $M \rightarrow N$ is automatically continuous.

Similarly, for two Banach K -spaces V and W we have that

$$\begin{aligned} \text{Hom}(V_0, W_0) &= \{f \in \text{Hom}(V, W) : f(V_0) \subseteq W_0\} \\ &= \text{Hom}_0(V, W) \end{aligned}$$

where the first equality follows as in the last sentence of the previous paragraph.

Finally, we observe that

$$(M \otimes_{\theta} K)_0 = M, \quad V_0 \otimes_{\theta} K = V$$

from where the proposition follows. \square

3.A.5 Banach spaces in terms of complete modules

We would now like to put this altogether to obtain Ban_K is a localization of Mod_θ^\wedge . Namely, let us define a morphism f in Mod_θ^\wedge to be a *topological isogeny* if $\ker f$ and $\text{coker}(f)$ have dense torsion submodules. We then have the following:

Proposition 3.A.10. *The functor*

$$F : \text{Mod}_\theta^\wedge \rightarrow \text{Ban}_K : M \mapsto (M/M_{\text{tors}})^\wedge \otimes_\theta K$$

realizes Ban_K as the localization of Mod_θ^\wedge at the set of topological isogenies.

Proof. By [?, Lemma 5.5] it suffices verify that F is essentially surjective, weakly full with fixed target (as in loc. cit.), and for all V in Ban_K we have that $F^{-1}(V)$ is a cofiltering category, and that $F(f)$ is an isomorphism if and only if f is topological isogeny.

To see that F is essentially surjective and weakly full with fixed target, we can apply Proposition 3.A.8.

To see that $F^{-1}(V)$ is cofiltering is clear

Finally, we verify that $F(f)$ is an isomorphism if and only if f is a topological isogeny. But, by the open mapping theorem we know that $F(f)$ is an isomorphism if and only if

$$\ker F(f) = \ker(f) \otimes_\theta K, \quad \text{coker}(F(f)) = \text{coker}(f) \otimes_\theta K$$

(using the θ -flatness of K) are both trivial. Thus, it suffices to show that $F(M)$ is zero if and only if M_{tors} is dense in M . But, since $(M/M_{\text{tors}})^\wedge$ is flat we know that $(M/M_{\text{tors}})^\wedge$ embeds into $F(M)$ and thus $F(M)$ is zero if and only if $(M/M_{\text{tors}})^\wedge = 0$. The claim then follows from Corollary 3.A.5. \square

4

Affinoid algebras and spaces

In this short section, we study quotients of Tate algebras, called *affinoid algebras*. The main result is that they carry natural equivalence classes of Banach K -algebra norms. Later, we will define their *affinoid spectra*, which will serve as building blocks for rigid-analytic spaces over K , just as spectra of finitely generated algebras over a field k are building blocks for schemes locally of finite type over k .

4.1 Affinoid algebras and the residue norm

Definition 4.1.1. Let K be a non-Archimedean field. A K -algebra A is an *affinoid algebra* if it is isomorphic to a quotient of the Tate algebra $K\langle X_1, \dots, X_r \rangle$ for some $r \geq 0$.

The results from §3.4 and §3.6 imply the following.

Proposition 4.1.2. *Every affinoid K -algebra A is Noetherian, Jacobson, and there exists a finite and injective K -algebra homomorphism*

$$K\langle Y_1, \dots, Y_s \rangle \hookrightarrow A$$

for some $s \geq 0$.

Let A be an affinoid K -algebra and let

$$\alpha: K\langle X_1, \dots, X_r \rangle \rightarrow A$$

be a surjective homomorphism; set $I = \ker(\alpha)$. Since every ideal in the Banach K -algebra $K\langle X_1, \dots, X_r \rangle$ is closed (Proposition 3.4.5), the quotient $A = K\langle X_1, \dots, X_r \rangle / I$ is a Banach space for the *residue norm*

$$|f|_\alpha = \inf\{|g| : g \in \alpha^{-1}(f)\}.$$

Further, it is trivial to check that $|\cdot|_\alpha$ is sub-multiplicative, therefore making $(A, |\cdot|_\alpha)$ into a Banach K -algebra. We shall soon prove that different presentations α give rise to *equivalent* norms $|\cdot|_\alpha$.

4.2 The supremum norm

Our goal in this section is to show that the K -algebra structure on an affinoid K -algebra A determines its topology. This is similar to the fact that the t -adic topology on an \mathcal{O} -module is canonically determined.

Our main foothold will be the corresponding result for finite field extensions of K , Theorem 2.5.1. We already know that affinoid K -algebras are Jacobson, which means that their maximal ideals carry significant information, and that the residue fields at maximal ideals are finite extensions of K . Together, these observations allow us to define the *supremum semi-norm* on an affinoid K -algebra A by setting

$$|f|_{\text{sup}} = \sup\{|f(x)| : x \in \text{Max}A\},$$

where $|f(x)|$ is the absolute value of the image of f in the residue field L of x with respect to the unique extension of the norm on K to L . (We already saw a preview of this for $A = K\langle X_1, \dots, X_r \rangle$ in the proof of Corollary 3.5.2.)

Proposition 4.2.1 (Properties of the supremum semi-norm). *Let A be an affinoid K -algebra.*

- (a) *The supremum semi-norm $|\cdot|_{\text{sup}}$ on A satisfies the axioms (i), (iii), and (iv) of a Banach K -algebra norm (Definition 3.2.5). It is power-multiplicative, in the sense that $|a^n|_{\text{sup}} = |a|_{\text{sup}}^n$. For every K -algebra homomorphism $\varphi: A \rightarrow B$ between affinoid algebras, we have $|\varphi(a)|_{\text{sup}} \leq |a|_{\text{sup}}$ for all $a \in A$.*
- (b) *One has $|a|_{\text{sup}} = 0$ if and only if a is nilpotent. If A is reduced, so that axiom (ii) of Definition 3.2.5 is also satisfied, then $|\cdot|_{\text{sup}}$ is a Banach K -algebra norm.*
- (c) *For $A = K\langle X_1, \dots, X_r \rangle$, the supremum norm coincides with the Gauss norm.*
- (d) *(Maximum principle) For every $a \in A$ there exists an $x \in \text{Max}A$ such that $|a|_{\text{sup}} = |a(x)|$. In particular, there exists an $n \geq 1$ such that $|a|_{\text{sup}}^n \in |K|$.*
- (e) *For every residue norm $|\cdot|_{\alpha}$ on A , an element $a \in A$ is powerbounded (Definition 3.3.1) if and only if $|a|_{\text{sup}} \leq 1$.*

Proof. Part (a) is clear. The first assertion of (b) follows from the fact that A is Jacobson, so that

$$\sqrt{(0)} = \bigcap_{\mathfrak{n} \in \text{Max}A} \mathfrak{n}.$$

Completeness of a reduced A with respect to $|\cdot|_{\text{sup}}$ is more involved and will not be needed; see [3, Theorem 6.2.4/1]. Part (c) was proved as part of the proof of Corollary 3.5.2.

For the remaining claims (d) and (e), we need some preparatory results. The following easy lemma says that one can estimate the absolute values of the roots of a polynomial by looking at its Newton polygon.

Lemma 4.2.2. *Let $f = X^n + a_1X^{n-1} + \dots + a_n \in K[X]$ be a polynomial, and let $\alpha_1, \dots, \alpha_n \in \overline{K}$ be its roots. Then*

$$\max_{i=1, \dots, n} |\alpha_i| = \max_{i=1, \dots, n} |a_i|^{1/i}.$$

Proof. The right-hand side is equal to $\exp(-\mu)$ where μ is the largest slope of $\text{NP}(f)$ (see Figure 4.2). By Lemma 2.6.2, this equals $\max |\alpha_i|$. \square




Figure 4.21: Proof of Lemma 4.2.2.

Let us fix a surjection $\alpha: K\langle X_1, \dots, X_r \rangle \rightarrow A$ and finite and injective homomorphism $\beta: K\langle Y_1, \dots, Y_s \rangle \rightarrow A$. By Remark 3.4.3, β can be lifted to a map $\gamma: K\langle Y_1, \dots, Y_s \rangle \rightarrow K\langle X_1, \dots, X_r \rangle$; part (c) implies that γ is contracting with respect to the Gauss norms.

We fix an $a \in A$; since a is integral over $K\langle Y_1, \dots, Y_s \rangle$, we fix a polynomial

$$f = X^n + f_1 X^{n-1} + \dots + f_n \in K\langle Y_1, \dots, Y_s \rangle[X]$$

such that $f(a) = 0$. We make the following assumption:¹

$$B = K\langle Y_1, \dots, Y_s \rangle[X]/(f) \rightarrow A \text{ is injective.}$$

Note that $B = K\langle Y_1, \dots, Y_s, X \rangle/(f)$ is also an affinoid K -algebra. Under the above assumption, $\text{Max}(A) \rightarrow \text{Max}(B)$ is surjective. Therefore

$$\begin{aligned} |a|_{\text{sup}} &= \sup_{x \in \text{Max}(A)} |a(x)| = \sup_{x \in \text{Max}(B)} |X(x)| \\ &= \sup_{y \in \text{Max}(K\langle Y_1, \dots, Y_s \rangle)} \max_{x \in \text{Max}(B), x \mapsto y} |X(x)|. \end{aligned}$$

By Lemma 4.2.2, the maximum equals $\max |f_i(y)|^{1/i}$, and hence the above equals $\max |f_i|_{\text{sup}}^{1/i} = \max |f_i|^{1/i}$.

We have thus, under our simplifying assumption, obtained the following assertion:

$$\text{One can find } f \text{ such that } |a|_{\text{sup}} = \max_{i=1, \dots, n} |f_i|^{1/i}.$$

We omit the rather unenlightening reduction to this case, referring the reader to [4, §3.1].²

To prove (d), we apply the Maximum Principle (Proposition 3.2.3) to $g = f_1 \cdot \dots \cdot f_n \in K\langle Y_1, \dots, Y_r \rangle$, obtaining a $y \in \text{Max} K\langle Y_1, \dots, Y_r \rangle$ with $|g|_{\text{sup}} = |g| = |g(y)|$. But this implies that $|f_i|_{\text{sup}} = |f_i| = |f_i(y)|$ for every i , and hence

$$|a|_{\text{sup}} = \max_{i=1, \dots, n} |f_i|^{1/i} = \max_{i=1, \dots, n} |f_i(y)|^{1/i} = \max_{x \mapsto y} |a(x)|.$$

To prove (e), the condition $|a|_{\text{sup}} \leq 1$ is equivalent to $|f_i| \leq 1$ for all i . This implies that a is integral over $\mathcal{O}\langle Y_1, \dots, Y_r \rangle$. Since γ is contracting (Corollary 3.5.2), the images $a_i = \beta(Y_i) = \alpha(\gamma(Y_i)) \in A$ satisfy $|a_i|_{\alpha} \leq 1$. This easily implies that a is power-bounded: if $C = \max\{|a^i|_{\alpha} : i < n\}$ then by induction we show that $|a^{n+m}|_{\alpha} \leq C$ for all $m \geq 0$:

$$|a^{n+m}|_{\alpha} = \left| -\sum_{i=0}^{n-1} a_{n-i} a^{i+m} \right|_{\alpha} \leq C.$$

Finally, if a is powerbounded, then $|a|_{\text{sup}}^n = |a^n|_{\text{sup}} \leq |a^n|_{\alpha}$ is bounded, forcing $|a|_{\text{sup}} \leq 1$. \square

Theorem 4.2.3. *Every K -algebra homomorphism $A \rightarrow B$ between affinoid K -algebras is continuous with respect to any choice of residue norms on the source and target. In particular, all residue norms on an affinoid K -algebra are equivalent.*

$$\begin{array}{ccc} K\langle X_1, \dots, X_r \rangle & \xrightarrow{\alpha, \text{ surjection}} & A \\ \uparrow \gamma, \text{ contracting} & \nearrow \beta, \text{ finite+injective} & \\ K\langle Y_1, \dots, Y_s \rangle & & \end{array}$$

¹ This assumption is satisfied for example if A is a domain, or just torsion-free as a $K\langle Y_1, \dots, Y_s \rangle$ -module, and if f is of minimal degree, see [4, Lemma 3.1/13].

² Idea of the reduction [4, Lemma 3.1/14]: replace A with $\prod A/\mathfrak{p}_i$ where $\mathfrak{p}_1, \dots, \mathfrak{p}_m \subseteq A$ are the minimal prime ideals. For each i , the affinoid K -algebra A/\mathfrak{p}_i is a domain, and we can apply [4, Lemma 3.1/13].

Proof. Fix a surjection $\alpha: K\langle X_1, \dots, X_r \rangle \rightarrow A$ corresponding to a residue norm $|\cdot|_\alpha$ and let

$$\varphi: K\langle X_1, \dots, X_r \rangle \rightarrow A \rightarrow B$$

be the composition. Since A has the quotient topology, it is enough to show that φ is continuous. In other words, we may replace A with $K\langle X_1, \dots, X_r \rangle$ endowed with the Gauss norm.

The elements $b_i = \varphi(X_i) \in B$ are power-bounded by Proposition 4.2.1(e), since

$$|b_i|_{\text{sup}} \leq |X_i|_{\text{sup}} = 1.$$

By the universal property of $K\langle X_1, \dots, X_r \rangle$ among Banach K -algebras, there exists a unique *continuous* K -algebra homomorphism

$$\varphi': K\langle X_1, \dots, X_r \rangle \rightarrow B \quad \text{such that} \quad \varphi'(X_i) = b_i.$$

It suffices to show that $\varphi = \varphi'$. Fix $f \in K\langle X_1, \dots, X_r \rangle$ and set $g = \varphi(f) - \varphi'(f) \in B$. For every maximal ideal $\mathfrak{n} \subseteq B$ and every $s \geq 1$, the quotient B/\mathfrak{n}^s is a finite dimensional K -algebra, and therefore the composition

$$\pi \circ \varphi: K\langle X_1, \dots, X_r \rangle \rightarrow B \rightarrow B/\mathfrak{n}^s$$

is continuous (since B/\mathfrak{n}^s is finite-dimensional, all norms are equivalent). Indeed, we may assume that $\pi \circ \varphi$ is surjective, and then $\pi \circ \varphi$ is continuous with respect to the residue norm it induces on B/\mathfrak{n}^s . This forces $\pi\varphi = \pi\varphi'$ by the universal property of $K\langle X_1, \dots, X_r \rangle$ applied this time to B/\mathfrak{n}^s .

Thus g maps to zero in B/\mathfrak{n}^s for every \mathfrak{n} and s . Therefore for every \mathfrak{n} , the image of g in $A_{\mathfrak{n}}$ lies in $\bigcap_s \mathfrak{n}^s A_{\mathfrak{n}}$, which is zero (by Krull's intersection theorem). This implies that $g = 0$. \square

4.3 The canonical topology

Let $A = K\langle X_1, \dots, X_r \rangle / (f_1, \dots, f_s)$ be an affinoid K -algebra. Then $\text{Max}A$ is identified with

$$\{x \in \mathbf{D}^r(\overline{K}) : f_i(x) = 0\} / \text{Gal}(\overline{K}/K).$$

Endowing \overline{K} with the metric topology, \overline{K}^r with the product topology, the set $\{f_i(x) = 0, |x_i| \leq 1\} \subseteq \overline{K}^r$ with the induced topology, and finally the above quotient by Galois action with the quotient topology, we obtain a topology on $\text{Max}A$ called the *canonical*. A more intrinsic (evidently independent of the presentation) definition is the following.

Definition 4.3.1. The *canonical topology* on $\text{Max}A$ is the topology generated by the subsets

$$X(f) = \{x \in \text{Max}A : |f(x)| \leq 1\}, \quad f \in A.$$

(More coming soon.)

Sheaves, sites, and topoi

5.1 Motivation: reinventing the real

Imagine being a geometer who does not believe in irrational numbers, perhaps for the fear of drowning. You study the geometry of the “line” \mathbf{Q} and maybe the higher-dimensional spaces \mathbf{Q}^r . With the irrationals hiding in your blind spot, the “unit interval” $[0, 1]_{\mathbf{Q}} = [0, 1] \cap \mathbf{Q}$ appears to you as both connected and compact, in the naive sense that it is not the union of two disjoint intervals *with rational endpoints*, and that every family of such intervals in \mathbf{Q} which covers $[0, 1]_{\mathbf{Q}}$ admits a finite subcover. Further, the functor assigning to each interval with rational endpoints $(a, b)_{\mathbf{Q}} = (a, b) \cap \mathbf{Q}$ the set of all continuous piecewise linear functions $(a, b)_{\mathbf{Q}} \rightarrow \mathbf{Q}$ satisfies the sheaf condition for *finite* coverings by intervals with rational endpoints.

Naturally, these properties fail to hold for the usual metric topology on \mathbf{Q} . Since we want to make do with what we have and avoid “filling in the holes,” we need a different way of formalizing our naive thoughts above.

Definition 5.1.1. A closed (resp. open) *rational box* is a subset of \mathbf{Q}^r of the form $\prod_{i=1}^r [a_i, b_i]_{\mathbf{Q}}$ (resp. $\prod_{i=1}^r (a_i, b_i)_{\mathbf{Q}}$) with $a_i, b_i \in \mathbf{Q}$.

- (a) An open subset $U \subseteq \mathbf{Q}^r$ is an *admissible open* if for every closed rational box $K \subseteq U$ there exists a finite collection V_1, \dots, V_m of open rational boxes contained in U such that $K \subseteq \bigcup_{i=1}^m V_i$.
- (b) A cover $U = \bigcup_{\alpha \in I} U_{\alpha}$ of an admissible open U by admissible opens U_{α} is an *admissible cover* if for every closed rational box $K \subseteq U$ there exists a finite collection V_1, \dots, V_m of open rational boxes contained in U such that $K \subseteq \bigcup_{i=1}^m V_i$ and each V_i is contained in some U_{α} .

Note that the intersection $U'' = U \cap U'$ of two admissible opens is again admissible. Indeed, if $K \subseteq U \cap U'$, we can find V_1, \dots, V_m and V'_1, \dots, V'_n as in the definition. Then $V''_{ij} = V_i \cap V'_j$ ($1 \leq i \leq m, 1 \leq j \leq n$) are rational boxes, are contained in U'' , and cover K .

Definition 5.1.2. A *sheaf for the admissible topology* on \mathbf{Q}^r is a functor¹

$$\mathcal{F}: \{\text{admissible opens in } \mathbf{Q}^r\} \rightarrow \mathbf{Sets}$$

such that for every admissible cover $U = \bigcup_{\alpha \in I} U_{\alpha}$ the sequence

$$\mathcal{F}(U) \rightarrow \prod_{\alpha \in I} \mathcal{F}(U_{\alpha}) \rightrightarrows \prod_{\alpha, \beta \in I} \mathcal{F}(U_{\alpha} \cap U_{\beta}) \quad (5.1)$$

Largely stolen from Brian Conrad’s lecture notes.

Convention: $(a, b)_{\mathbf{Q}} = (a, b) \cap \mathbf{Q}$ etc.

¹ Here we regard any poset C as a category with morphisms

$$\mathrm{Hom}_C(c, c') = \begin{cases} \{*\} & \text{if } c \leq c' \\ \emptyset & \text{otherwise.} \end{cases}$$

is exact.²

Recall some basic terminology: if $U = \bigcup_{\alpha \in I} U_\alpha$ is an open cover, we say that \mathcal{F} satisfies the sheaf condition for $\{U_\alpha\}_{\alpha \in I}$ if (5.1) is exact. If $U = \bigcup_{\beta \in J} V_\beta$ is another cover, we say that $\{V_\beta\}_{\beta \in J}$ refines $\{U_\alpha\}_{\alpha \in I}$ if every V_β is contained in some U_α ; more precisely, if there exists a map $f: J \rightarrow I$ such that $V_\beta \subseteq U_{f(\beta)}$ for every $\beta \in J$.

Proposition 5.1.3. (a) Let \mathcal{G} be a functor from the poset of closed rational boxes in \mathbf{Q}^r to sets which satisfies the sheaf condition for every finite covering $K = \bigcup_{\alpha \in I} K_\alpha$. Then \mathcal{G} extends uniquely to a sheaf for the admissible topology on \mathbf{Q}^r .

(b) If $\overline{\mathcal{F}}$ is a sheaf on \mathbf{R}^r (for the standard topology), then the functor associating to a closed rational box $K = \prod [a_i, b_i]_{\mathbf{Q}}$ the value

$$\overline{\mathcal{F}}(\prod [a_i, b_i]) := \varinjlim_{\prod [a_i, b_i] \subseteq U \subseteq \mathbf{R}^r} \overline{\mathcal{F}}(U)$$

satisfies the sheaf condition for every finite covering of a closed rational box by closed rational boxes, and therefore by (a) it extends uniquely to a sheaf for the admissible topology on \mathbf{Q}^r , denoted \mathcal{F} .

(c) The association $\overline{\mathcal{F}} \mapsto \mathcal{F}$ defines an equivalence of categories

$$\{\text{sheaves on } \mathbf{R}^r\} \simeq \{\text{sheaves for the admissible topology on } \mathbf{Q}^r\}.$$

Proof. Omitted, but see Problems 2 and 3 on Problem Set 4. □

In Appendix 5.A we will learn how to reconstruct certain topological spaces from their category of sheaves. In particular, we shall obtain:

Corollary 5.1.4 (See Appendix 5.A). *The space \mathbf{R}^r can be recovered from the category of sheaves for the admissible topology on \mathbf{Q}^r .*

Example 5.1.5. (a) Every open subset $U \subseteq \mathbf{Q}$ is admissible.

(b) However, the covering of \mathbf{Q} by all open rational intervals $(a, b)_{\mathbf{Q}}$ such that $\sqrt{2} \notin (a, b)$ is not an admissible cover, since e.g. $K = [0, 1]_{\mathbf{Q}}$ cannot be covered by finitely many such intervals.

(c) The sheaf “skyscraper at $\sqrt{2}$,” defined as

$$\overline{\mathcal{F}}(U) = \begin{cases} \mathbf{Z} & \text{if } \sqrt{2} \in U \\ 0 & \text{otherwise} \end{cases}$$

defines a nonzero sheaf \mathcal{F} for the admissible topology on \mathbf{Q} whose stalks at all points in \mathbf{Q} (defined in the obvious way) are zero.

(d) The following is an example of an inadmissible open in \mathbf{Q}^2 (due to Zev Rosengarten):

$$U = \mathbf{Q}^2 \cap \left((0, \sqrt{2}) + \{x \geq -|y|\} \right) \quad (\text{see Figure 5.1}).$$

In this case, the closed box $K = [0, 1] \times [0, 2]$ does not admit a finite cover by open subsets contained in U .

² Here **exact** is another name for an *equalizer*: the left map is injective and its image equals the set of elements whose images by the two parallel arrows are equal.

The consideration of values on closed boxes is a bit artificial here. In algebraic geometry and rigid geometry, our basic opens (affine or affinoid opens) will be quasi-compact, and there will be no need to consider the values of a sheaf on closed sets.




Figure 5.11: An inadmissible open subset of \mathbf{Q}^2 .

5.2 Sites

Definition 5.2.1 (Site). A *site* is a category \mathcal{C} in which every object $c \in \text{ob } \mathcal{C}$ is endowed with a collection $\text{Cov } c$ of families of maps $\{c_\alpha \rightarrow c\}_{\alpha \in I}$, called *covering families*, satisfying the following axioms.

- i. (ISOMORPHISM) If $c' \rightarrow c$ is an isomorphism then the singleton $\{c' \rightarrow c\}$ is a covering family of c ,
- ii. (PULLBACK) If $\{c_\alpha \rightarrow c\}_{\alpha \in I}$ is a covering family of c and if $c' \rightarrow c$ is a morphism, then the fiber products $c'_\alpha = c_\alpha \times_c c'$ exist and the family

$$\{c'_\alpha = c_\alpha \times_c c' \rightarrow c'\}_{\alpha \in I}$$

is a covering family of c' .

- iii. (COMPOSITION) If $\{c_\alpha \rightarrow c\}_{\alpha \in I}$ is a covering family of c and for every $\alpha \in I$ we have a covering family $\{c_{\alpha\beta} \rightarrow c_\alpha\}_{\beta \in I_\alpha}$ of c_α , then

$$\{c_{\alpha\beta} \rightarrow c_\alpha \rightarrow c\}_{\alpha \in I, \beta \in I_\alpha}$$

is a covering family of c .

The basic example is of course the site $\text{Op } X$ of opens in a topological space X , where morphisms are inclusions $U' \subseteq U$ of open subsets, and where $\{U_\alpha \subseteq U\}$ is a covering family precisely when $U = \bigcup U_\alpha$. Another one is provided by our toy example above: the category of admissible opens in \mathbf{Q}^r where covering families are given by admissible covers.³

Note that axioms (i) and (ii) imply that an isomorphism $c' \rightarrow c$ induces a bijection between covering families of c and of c' . By abuse of terminology, we shall use the notation \mathcal{C} to refer to both the site and the underlying category. A safer way would be to give a name such as τ to the choice of $\text{Cov } c$ for every $c \in \mathcal{C}$ satisfying the above axioms (called a *Grothendieck (pre)topology* on the category \mathcal{C}) and define a site as a category \mathcal{C} with a Grothendieck topology τ , denoted (\mathcal{C}, τ) . This is sometimes useful, e.g. if one considers two sites with the same underlying category.⁴

Definition 5.2.2 (Sheaf). Let \mathcal{C} be a site. A *sheaf* on \mathcal{C} is a contravariant functor

$$\mathcal{F} : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Sets}$$

such that for every $c \in \text{ob } \mathcal{C}$ and every covering family $\{c_\alpha \rightarrow c\}_{\alpha \in I}$ the sequence

$$\mathcal{F}(c) \rightarrow \prod_{\alpha \in I} \mathcal{F}(c_\alpha) \rightrightarrows \prod_{\alpha, \beta \in I} \mathcal{F}(c_\alpha \times_c c_\beta) \quad (5.2)$$

is exact (note that the fiber products $c_\alpha \times_c c_\beta$ exist thanks to axiom ii).

We denote by $\text{Sh } \mathcal{C}$ the category of sheaves on \mathcal{C} , considered as a full subcategory of the category of *presheaves* $\text{PSh } \mathcal{C} = \text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Sets})$. We call $\text{Sh } \mathcal{C}$ the *topos* associated to \mathcal{C} .

In general, a *topos* (plural: *topoi*) is a category which is equivalent to $\text{Sh } \mathcal{C}$ for some site \mathcal{C} (with no extra structure!). Different sites can give rise to equivalent topoi, and so a topos is in a way a superior notion; we

³ More examples of sites:

- The *étale site* of a scheme X : the objects are étale morphisms $U \rightarrow X$, maps are morphisms over X , and covers $\{U_\alpha \rightarrow U\}$ are families of jointly surjective maps,
- Replacing *étale* with *flat and locally finitely presented* one obtains the *fppf site*.
- For a group G , the category of G -sets where covers are jointly surjective families of G -equivariant maps.

⁴ The same happens in topology: one uses a letter such as X to denote both a topological space and the underlying set; if confusion is possible, one writes (X, \mathcal{T}) for the topological space.

can regard a site as a particular presentation of the associated topos, just as a metric on a topological space is a useful but non-canonical “presentation” of its topology.

So far, to define sheaves and topoi, we only needed a part of axiom (ii), namely that suitable fiber products exist. To see the other axioms in action, let us show that familiar features of sheaf theory: refinement, (zeroth) Čech cohomology, and sheafification, work in a similar way in a site \mathcal{C} .

Definition 5.2.3 (Refinement). We say that a covering family $\{c'_\beta \rightarrow c\}_{\beta \in I}$ refines a covering family $\{c_\alpha \rightarrow c\}_{\alpha \in I}$ of the same object c if there exists a function $\varphi: J \rightarrow I$ and maps $\varphi_\beta: c'_\beta \rightarrow c_{\varphi(\beta)}$ fitting inside a commutative triangle

$$\begin{array}{ccc} c'_\beta & \xrightarrow{\varphi_\beta} & c_{\varphi(\beta)} \\ & \searrow & \swarrow \\ & c & \end{array}$$

This is an analog of the usual notion in topology: a cover $U = \bigcup U'_\beta$ refines $U = \bigcup U_\alpha$ if every U'_β is contained in some U_α . The relation of refinement is clearly transitive. Further, axioms (ii) and (iii) imply that every two covering families $\{c_\alpha \rightarrow c\}_{\alpha \in I}$ and $\{c'_\beta \rightarrow c\}_{\beta \in J}$ admit a common refinement, namely

$$\{c_\alpha \times_c c'_\beta \rightarrow c\}_{(\alpha, \beta) \in I \times J}.$$

Given a presheaf $\mathcal{F}: \mathcal{C}^{\text{op}} \rightarrow \mathbf{Sets}$ and a covering family $\{c_\alpha \rightarrow c\}_{\alpha \in I}$ let us define $\mathcal{H}^0(\mathcal{F}, \{c_\alpha\})$ as the equalizer of

$$\prod_{\alpha \in I} \mathcal{F}(c_\alpha) \rightrightarrows \prod_{\alpha, \beta \in I} \mathcal{F}(c_\alpha \times_c c_\beta).$$

Thus \mathcal{F} satisfies the sheaf condition for $\{c_\alpha\}$ (meaning that (5.2) is exact) precisely when the canonical map $\mathcal{F}(c) \rightarrow \mathcal{H}^0(\mathcal{F}, \{c_\alpha\})$ is a bijection.

Lemma 5.2.4. *Let \mathcal{F} be a presheaf on \mathcal{C} and let $\{c_\alpha \rightarrow c\}, \{c'_\beta \rightarrow c\}$ be two covering families of an object c .*

(a) *Let $\varphi: J \rightarrow I$ and $\varphi_\beta: c'_\beta \rightarrow c_{\varphi(\beta)}$ be as in Definition 5.2.3. Then $(\varphi, \{\varphi_\beta\})$ induces a map*

$$\mathcal{H}^0(\mathcal{F}, \{c_\alpha\}) \rightarrow \mathcal{H}^0(\mathcal{F}, \{c'_\beta\}).$$

(b) *If $\varphi': J \rightarrow I$, $\varphi'_\beta: c'_\beta \rightarrow c_{\varphi'(\beta)}$ is another such datum, then the two induced maps $\mathcal{H}^0(\mathcal{F}, \{c_\alpha\}) \rightarrow \mathcal{H}^0(\mathcal{F}, \{c'_\beta\})$ are equal.*

This means that if $\{c'_\beta \rightarrow c\}$ refines $\{c_\alpha \rightarrow c\}$, we obtain a canonical map $\mathcal{H}^0(\mathcal{F}, \{c_\alpha\}) \rightarrow \mathcal{H}^0(\mathcal{F}, \{c'_\beta\})$. Thus if we consider $\text{Cov } c$ as a partially ordered set with respect to the relation of refinement (as we observed, this poset is cofiltering: every two elements have a common upper bound), we can define the *zeroth Čech cohomology* as the colimit

$$\check{\mathcal{H}}^0(\mathcal{F}, c) = \varinjlim_{\{c_\alpha\} \in \text{Cov } c} \mathcal{H}^0(\mathcal{F}, \{c_\alpha\}).$$

Then $c \mapsto \check{\mathcal{H}}^0(\mathcal{F}, c)$ is another presheaf on \mathcal{C} , denoted \mathcal{F}^+ .

[10, Tag 00W6]

[10, Tag 00W7]

Lemma 5.2.5 (Sheafification). *For every presheaf \mathcal{F} on \mathcal{C} , the presheaf $(\mathcal{F}^+)^+$ is a sheaf. The functor $\mathcal{F} \mapsto \mathcal{F}^\# := (\mathcal{F}^+)^+$ is a left adjoint to the inclusion $\text{Sh } \mathcal{C} \subseteq \text{PSh } \mathcal{C}$, called the sheafification functor.*

Further, many other notions of sheaf theory: cohomology, continuous maps of sites $f: \mathcal{C} \rightarrow \mathcal{C}'^5$, push-forward and pull-back functors $f_*: \text{Sh } \mathcal{C} \rightarrow \text{Sh } \mathcal{C}'$ and $f^*: \text{Sh } \mathcal{C}' \rightarrow \text{Sh } \mathcal{C}$, and so on, exist and behave as one would expect.

Let us stop here the development of the general theory, referring the curious reader to [12], [9], [1], or [10, Tag 00UZ].

5.3 G -topologies

The admissible site of \mathbf{Q}^r defined in §5.1 is fairly concrete: its objects are simply subsets of the set \mathbf{Q}^r . In other words, admissible opens and covers define a G -topology in the sense of the following definition.

Definition 5.3.1 (G -topology). A G -topology on a set X is a site whose underlying category is a full subcategory of the poset of subsets of X , which is stable under intersections and such that covering families are jointly surjective.

In other words, it is the data of a set \mathcal{C} of subsets of X , called *admissible opens*, such that the intersection of two admissible subsets is again admissible, and for each admissible open $U \in \mathcal{C}$, a class of *admissible covers* $\{U_\alpha\}_{\alpha \in I}$ where $U = \bigcup_{\alpha \in I} U_\alpha$ and $U_\alpha \in \mathcal{C}$, such that the following axioms are satisfied

- i. The cover $\{U\}$ is an admissible cover for every $U \in \mathcal{C}$.
- ii. If $U' \subseteq U$ is an inclusion of admissible opens and if $\{U_\alpha\}$ is an admissible cover of U , then $\{U'_\alpha = U_\alpha \cap U'\}$ is an admissible cover of U' .
- iii. If $\{U_\alpha\}_{\alpha \in I}$ is an admissible cover of an admissible open U and if for every $\alpha \in I$, $\{U_{\alpha\beta}\}_{\beta \in J_\alpha}$ is an admissible cover of U_α , then $\{U_{\alpha\beta}\}_{\alpha \in I, \beta \in J_\alpha}$ is an admissible cover of U .

A G -topological space is a set X endowed with a G -topology.

Example 5.3.2. Let X be a separated scheme, and take as admissible opens the set of all affine open subsets $U \subseteq X$. Separatedness ensures that \mathbf{C} is stable under pairwise intersection. There are two variants of admissible covers:

- (STRONG) A covering $\{U_\alpha\}_{\alpha \in I}$ of an affine open U by affine opens U_α is admissible if $U = \bigcup U_\alpha$.
- (WEAK) The same but with I finite.

Since every affine scheme is quasi-compact, both give rise to the same category of sheaves, which is moreover equivalent to the category of sheaves on the topological space X .

[10, Tag 00WB], [10, Tag 00WH]

⁵ By convention, this is a functor $f^{-1}: \mathcal{C}' \rightarrow \mathcal{C}$ in the opposite direction. It is assumed to map covering families to covering families and to preserve fiber products.

These conditions ensure that the functor

$$(-) \circ f^{-1}: \text{PSh } \mathcal{C} \rightarrow \text{PSh } \mathcal{C}'$$

maps sheaves to sheaves, inducing a functor

$$f_*: \text{Sh } \mathcal{C} \rightarrow \text{Sh } \mathcal{C}'.$$

In a G -topological space, we say that some property holds *locally* if it does so on the members of an admissible covering.

Our goal in the next chapter will be to put a G -topology on the space of maximal ideals of an affinoid K -algebra, as well as a structure sheaf. We will then glue such spaces to obtain general rigid-analytic spaces. Gluing G -topologies is facilitated by the following properties:

Definition 5.3.3 (Completeness axioms). Let X be a G -topological space.

- (0) We say that X satisfies axiom (G_0) if \emptyset and X are admissible opens.
- (1) We say that X satisfies axiom (G_1) if “admissibility of a subset is a local condition”: given a subset $V \subseteq U$ of an admissible open U , the set V is an admissible open if and only if there exists an admissible cover $\{U_\alpha\}$ of U such that $U_\alpha \cap V$ is an admissible open for all α .
- (2) We say that X satisfies axiom (G_2) if a covering of an admissible open V by admissible opens $\{V_\alpha\}$ is admissible if it admits an admissible covering of V as a refinement.

Remark 5.3.4. Consider the following condition (G'_2) “admissibility of a cover is a local condition”: given an admissible open V contained in an admissible open U and a family $\{V_\beta\}$ of admissible open subsets of V , the family $\{V_\beta\}$ is an admissible cover of V if and only if there exists an admissible cover $\{U_\alpha\}$ of U such that $\{U_\alpha \cap V_\beta\}$ is an admissible cover of $U_\alpha \cap V$ for every α . Then $(G_2) \Rightarrow (G'_2)$, and if the G -topology satisfies the additional property that every cover of the form $V = \bigcup V_\alpha$ of an admissible open by admissible opens such that $V = V_\alpha$ for some α is admissible (that is, “split” covers are admissible), then also $(G'_2) \Rightarrow (G_2)$.

If U is an admissible open of a G -topological space X , then the set of all admissible opens $V \subseteq U$ and the datum of all admissible covers consisting of such subsets forms a G -topology on U , called the *induced G -topology*.

[4, Proposition 5.1/11]

Proposition 5.3.5 (Gluing G -topologies). Let X be a set and let $U_\alpha \subseteq X$ ($\alpha \in I$) be subsets of X such that $X = \bigcup U_\alpha$. Suppose that

- each U_α is endowed with a G -topology satisfying axioms (G_0) , (G_1) , and (G_2) , and
- $U_\alpha \cap U_\beta$ is an admissible open in both U_α and U_β for every $\alpha, \beta \in I$, and
- the G -topologies on U_α and U_β induce the same G -topology on $U_\alpha \cap U_\beta$.

Then there exists a unique G -topology on X satisfying (G_0) , (G_1) , and (G_2) for which the U_α are admissible opens, for which $X = \bigcup U_\alpha$ is an admissible cover, and which induces the given topology on each U_α .

Proof. Condition (G_1) imposes that $V \subseteq X$ is admissible if and only if $V \cap U_\alpha$ is an admissible open of U_α for all α . Similarly, (G_2) forces declaring $\{V_\beta\}$ an admissible cover of $V = \bigcup V_\beta$ if $\{U_\alpha \cap V_\beta\}_\beta$ is an admissible cover of $U_\alpha \cap V$ (for the G -topology on U_α) for every α . This shows uniqueness, and we need to check that this defines a G -topology on X with the desired properties. This is rather straightforward and we omit the proof. \square

5.A Sober topological spaces

For a topological space X , let $\text{Op}(X)$ be the poset of open subsets of X , ordered by inclusion.⁶ Can we recover X from $\text{Op}(X)$? Clearly not always, for example if X has the indiscrete topology (the only opens are X and \emptyset) then $\text{Op}(X)$ carries no information about the cardinality of X . More generally, if X is not T_0 , i.e. there exist two points $x \neq x'$ which such that $x \in U \iff x' \in U$ for every open $U \subseteq X$, then $\text{Op}(X)$ and $\text{Op}(X/(x \sim x'))$ are isomorphic.

Even axiom T_0 is not sufficient for the recovery of X for $\text{Op}(X)$. For example, if $X = \mathbf{A}_k^1$ with the Zariski topology and $X' = X \setminus \{\eta\}$ is the set of all closed points of X (η is the generic point), then $\text{Op}(X) \simeq \text{Op}(X')$, since a non-empty $U \subseteq X'$ is open if and only if $U \cup \{\eta\}$ is open in X' .

Recall that a closed subset $Y \subseteq X$ of a topological space X is *irreducible* if it is not the sum of two proper closed subsets. If $Y = \overline{\{y\}}$ for some point $y \in Y$, we call y a *generic point* of Y .

Definition 5.A.1 (Sober space). A topological space X is *sober* if every irreducible closed subset $Y \subseteq X$ has a unique generic point. We denote by $\mathbf{Top}^{\text{sober}} \subseteq \mathbf{Top}$ the full subcategory of sober spaces.

Proposition 5.A.2. *The inclusion functor $\mathbf{Top}^{\text{sober}} \subseteq \mathbf{Top}$ admits a left adjoint $X \mapsto X^{\text{sob}}$, the soberification.*

Proof sketch. Let X be a topological space and let X^{sob} be the set of all irreducible closed subsets of X ; we have a natural map $\tau_X: X \rightarrow X^{\text{sob}}$ sending x to $\overline{\{x\}}$. We endow X^{sob} with the topology in which a subset $U \subseteq X^{\text{sob}}$ open if there exists an open $U^\circ \subseteq X$ such that U equals the set of irreducible subsets which intersect U° . This topology makes $\tau_X: X \rightarrow X^{\text{sob}}$ continuous. Moreover, the open subset U° is unique if it exists, so we have an order-preserving bijection $U \leftrightarrow U^\circ$ between opens in X and in X^{sob} .

The space X^{sob} is sober: if $Z \subseteq X^{\text{sob}}$ is an irreducible closed subset, write its complement $U = X^{\text{sob}} \setminus Z$ as the set of all closed irreducible $Y \subseteq X$ which intersect some open $U^\circ \subseteq X$. Set $W = X \setminus U^\circ$; it is easy to check that W is irreducible, and hence defines a point $[W] \in X^{\text{sob}}$. One then checks that $Z = \overline{[W]}$, and that $[W]$ is the unique generic point of Z . Details omitted.

If $f: X \rightarrow X'$ is continuous, and $Y \subseteq X$ is closed and irreducible, then $f(Y) \subseteq X'$ is irreducible, and so is its closure $\overline{f(Y)}$. The map $f^{\text{sob}}: X^{\text{sob}} \rightarrow (X')^{\text{sob}}$ defined by $Y \mapsto \overline{f(Y)}$ is continuous. Moreover, the square

$$\begin{array}{ccc} X & \xrightarrow{f} & X' \\ \tau_X \downarrow & & \downarrow \tau_{X'} \\ X^{\text{sob}} & \xrightarrow{f^{\text{sob}}} & (X')^{\text{sob}} \end{array}$$

commutes. We have thus defined a functor $X \mapsto X^{\text{sob}}: \mathbf{Top} \rightarrow \mathbf{Top}^{\text{sober}}$ and a natural transformation τ which will serve as the unit of the adjunction.

⁶ The poset $\text{Op}(X)$ is an example of a **locale**: a poset in which the supremum of every subset and the infimum of every finite subset exists, and which satisfies the *distributive law*

$$\inf\{x, \sup\{a_i\}_{i \in I}\} = \sup\{\inf\{x, a_i\}\}_{i \in I}.$$

Every locale forms a site where $\{a_i \leq a\}_{a \in A}$ forms a covering family if $a = \sup\{a_i\}$, and hence gives rise to a topos. For the locale $\text{Op}(X)$, this is the usual sheaf theory on X .

See [6, 0 2.1.(b)]

Finally, we need to check that every map $X \rightarrow X'$ with X' sober factors uniquely through X^{sob} . This is equivalent to saying that $\tau_X : X \rightarrow X^{\text{sob}}$ is a homeomorphism of X is sober. The inverse maps $[Y]$ to the unique generic point η_Y of Y ; it is clearly an inverse bijection. It is also continuous, since the preimage of $\{[Y] : Y \cap U^\circ \neq \emptyset\}$ equals U° . \square

Since $\tau_X : X \rightarrow X^{\text{sob}}$ induces a bijection on open subsets, we have $\text{Op}(X) \simeq \text{Op}(X^{\text{sob}})$ as posets. Conversely, the construction of the space X^{sob} only depends on the poset $\text{Op}(X)$. Indeed, the set of closed irreducible subsets Y of X is in bijection $Y \leftrightarrow X \setminus Y = U$ with the set of open subsets $U \in \text{Op}(X)$ which are not equal to the intersection $U_1 \cap U_2$ of two opens $U_1, U_2 \neq U$. Since $U_1 \cap U_2$ is the largest element of the poset $\text{Op}(X)$ which is smaller than both U_1 and U_2 , the latter depends only on the order on $\text{Op}(X)$. Summarizing:

Corollary 5.A.3. *The soberification of a space X depends only on the poset $\text{Op}(X)$, and the poset $\text{Op}(X)$ depends only on the soberification of X . For two spaces X and Y , there exists an isomorphism of posets $\text{Op}(X) \simeq \text{Op}(Y)$ if and only if $X^{\text{sob}} \simeq Y^{\text{sob}}$.*

For a family $\{U_\alpha\}$ of open subsets of a space X , the union $U = \bigcup U_\alpha$ is the smallest element of the poset $\text{Op}(X)$ which is larger than all U_α . It follows that the topos $\text{Sh}(X)$ (the category of sheaves on X) depends only on the poset $\text{Op}(X)$. In particular, X and X^{sob} have equivalent topoi.

It turns out that $\text{Op}(X) \mapsto \text{Sh}(X)$ does not lose any information, namely:

Proposition 5.A.4. *Let X be a sober topological space. Then X can be reconstructed from the topos $\text{Sh}(X)$.*

See [6, 0 2.7(a)].

Proof. Note that every topos $T = \text{Sh } \mathcal{C}$ admits a final object e , the sheaf whose value on every $c \in \text{ob } \mathcal{C}$ is the singleton $\{*\}$. If $T = \text{Sh}(X)$ for a topological space X , then $e = \text{Hom}_{\text{Op}(X)}(-, X)$ is simply the sheaf represented by X , the final object of the site $\text{Op}(X)$.

Ignoring potential set-theoretic difficulties, let us consider the set $\text{Op}(T)$ of sub-objects of e , i.e. isomorphism classes of objects $v \in \text{ob } T$ such that the unique morphism $v \rightarrow e$ is a monomorphism. We endow $\text{Op}(T)$ with the order where we declare $v \leq v'$ if there exists a morphism $v \rightarrow v'$.

Suppose now that $T = \text{Sh}(X)$ for a topological space X . If $U \subseteq X$ is an open subset, then the sheaf $h_U = \text{Hom}_{\text{Op}(X)}(-, U)$ is a sub-object of the final object h_X ; moreover, if $V \subseteq U$ then $h_V \leq h_U$, so we get a morphism of posets

$$\gamma : \text{Op}(X) \rightarrow \text{Op}(T).$$

We claim that γ is an isomorphism of posets. Indeed, if v is a sub-object of e , let U be the union of all opens $V \subseteq X$ such that $v(V) \neq \emptyset$. Since every $v(V)$ is a subset of $\{*\} = e(V)$, the sheaf condition implies that $v(U) = \{*\}$. By Yoneda's lemma, this gives a map of sheaves $h_U \rightarrow v$. This map is an isomorphism on stalks and hence is an isomorphism. This gives the inverse to γ , and we omit checking all the remaining details.

Finally, $X^{\text{sob}} = X$ can be reconstructed from $\text{Op}(X) \simeq \text{Op}(\text{Sh}(X))$ by Corollary 5.A.3. \square

We come back to our toy example at the beginning of the chapter:

Corollary 5.A.5. *The space \mathbf{R}^r can be reconstructed from the category $\mathrm{Sh}^{\mathrm{adm}}(\mathbf{Q}^r)$ of sheaves for the admissible topology on \mathbf{Q}^r .*

The same idea in rigid geometry recovers the *adic spectrum* $\mathrm{Spa}A$ of an affinoid K -algebra A in the sense of Huber from the affinoid space $(\mathrm{Sp}A = \mathrm{Max}A, \text{admissible topology})$, to be defined next. Thus a good understanding of the points of $\mathrm{Spa}A$ allows one to get rid of the G -topology in favor of usual topology.

6

Bibliography

- [1] *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos.* Lecture Notes in Mathematics, Vol. 269. Springer-Verlag, Berlin-New York, 1972. Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck, et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat.
- [2] M. F. Atiyah and I. G. Macdonald. *Introduction to commutative algebra.* Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [3] S. Bosch, U. Güntzer, and R. Remmert. *Non-Archimedean analysis*, volume 261 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1984. A systematic approach to rigid analytic geometry.
- [4] Siegfried Bosch. *Lectures on formal and rigid geometry*, volume 2105 of *Lecture Notes in Mathematics*. Springer, Cham, 2014.
- [5] Brian Conrad. Irreducible components of rigid spaces. *Ann. Inst. Fourier (Grenoble)*, 49(2):473–541, 1999.
- [6] Kazuhiro Fujiwara and Fumiharu Kato. *Foundations of rigid geometry. I.* EMS Monographs in Mathematics. European Mathematical Society (EMS), Zürich, 2018.
- [7] Robin Hartshorne. *Algebraic geometry.* Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [8] Jürgen Neukirch. *Algebraic number theory*, volume 322 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, 1999.
- [9] Martin Olsson. *Algebraic spaces and stacks*, volume 62 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2016.
- [10] The Stacks Project Authors. *Stacks Project*. <https://stacks.math.columbia.edu>, 2018.
- [11] Yichao Tian. Introduction to rigid geometry. online lecture notes.

- [12] Angelo Vistoli. Grothendieck topologies, fibered categories and descent theory. In *Fundamental algebraic geometry*, volume 123 of *Math. Surveys Monogr.*, pages 1–104. Amer. Math. Soc., Providence, RI, 2005.